

Observation of new resonances decaying to $J/\psi K^+$ and $J/\psi \phi$



张黎明 (清华大学)
2021年3月12日



强子物理
在线论坛






Z_c^- and earlier predictions of Z_{CS}

- Several Z_c^- states were observed from $Y(4260)$ or B decays, **at least have $c\bar{c}d\bar{u}$ four quarks**
- Would be nice to look for Z_{CS} , the SU(3) partners of $X(3872)/Z_c(3900)$
- It's useful to distinguish different models
 - Less exchange particles expected the Z_{CS} molecule picture
- Several papers have predicted the existence of Z_{CS} state in early time

[Phys. Soc. 55(2009)424, PRL 110(2013)232001, PRD 88(2013)096014, PLB 798(2019)135022, JHEP 04(2020)119]


Others from B decays are broad



In Nov. 2016 at XYZ workshop in Beihang, Xiaorui and I exchanged ideas to search for Z_{CS}


What triggered this analysis

- We first looked at $B^+ \rightarrow J/\psi\phi K^+$ in winter of 2019, as a control channel to study Cabibbo-suppressed $B^+ \rightarrow J/\psi\bar{K}^{*0}K^+$ motivated by Z_{CS} search.
- Instead, the control channel showed a possible $J/\psi K^+$ structure in Dalitz plot
- But at that time:
 - No exotic state with strangeness observed
 - The structure is not super narrow, amplitude analysis required
- We then prepared simulation sample, read the Run-1 amplitude analysis note, searched theoretical predictions about exotic states containing strange quark, tried to find other channels to confirm it ...



BESIII observation

- BESIII recently observed a **narrow** $Z_{cS}(3985)^-$ in $D_S^- D^{*+} + DD_S^{*-}$ mass
- Theory interpretations
 - Molecular partner of $Z_c(3900)^-$ from $D_S^- D^{*0} + D_S^{*-} D^0 +$ others exchanging $\eta/\sigma/f_0, 2K, c\bar{c}$
 - Diquark-antidiquark compact type
 - Kinematic reflection
- Some theorists points out 1^{++} and 1^{+-} may both exist in both Molecular and compact pictures [2011.10495,2011.10959,2012.11869]





System	Current	J^P	$m_H(\text{GeV})$
$\bar{D}_s D$	J_1	0^+	3.74 ± 0.13
$\bar{D}_s^* D^*$	J_2	0^+	4.11 ± 0.14
$\bar{D}_s^* D$	$J_{1\mu}$	1^+	3.99 ± 0.12
$\bar{D}_s D^*$	$J_{2\mu}$	1^+	3.97 ± 0.11
$\bar{D}_s^* D^*$	$J_{3\mu}$	1^+	4.22 ± 0.14
$\bar{D}_s^* D^*$	$J_{4\mu}$	1^+	4.22 ± 0.14
$\bar{D}_s^* D^*$	$J_{\mu\nu}$	2^+	4.34 ± 0.13
$\mathbf{0}_{[sc]} \oplus \mathbf{0}_{[\bar{q}\bar{e}]}$ (spin-spin)	η_1	0^+	3.84 ± 0.15
$\mathbf{1}_{[sc]} \oplus \mathbf{1}_{[\bar{q}\bar{e}]}$	η_2	0^+	4.13 ± 0.17
$\mathbf{1}_{[sc]} \oplus \mathbf{0}_{[\bar{q}\bar{e}]}$	$\eta_{1\mu}$	1^+	3.98 ± 0.16
$\mathbf{0}_{[sc]} \oplus \mathbf{1}_{[\bar{q}\bar{e}]}$	$\eta_{2\mu}$	1^+	3.97 ± 0.15
$\mathbf{1}_{[sc]} \oplus \mathbf{1}_{[\bar{q}\bar{e}]}$	$\eta_{3\mu}$	1^+	4.28 ± 0.14
$\mathbf{1}_{[sc]} \oplus \mathbf{1}_{[\bar{q}\bar{e}]}$	$\eta_{4\mu}$	1^+	4.28 ± 0.14
$\mathbf{1}_{[sc]} \oplus \mathbf{1}_{[\bar{q}\bar{e}]}$	$\eta_{\mu\nu}$	2^+	4.33 ± 0.13




Search for Z_{cs} at LHCb

- In Run 1 analysis, a hint of excess near 4 GeV, but not significant.
- Z_{cs}^+ in $B^+ \rightarrow J/\psi\phi K^+$ decay has similar topology as Z_c^- in $B^0 \rightarrow J/\psi K^+\pi^-$ decay, and P_c^+ in $\Lambda_b^0 \rightarrow J/\psi K^- p$, where spectator quark in b -hadron contributes to the exotic valence quarks.

[PRL118(2017)022003]





[PLB 798 (2019) 135022]



$X(4140)$ and $X(4274)$




- $B^+ \rightarrow J/\psi\phi K^+$ decays provide rich exotic spectra, initially used for study of $J/\psi\phi$ structures
- CDF observed a narrow $J/\psi\phi$ structure in [Initial publication on 2.7 fb⁻¹ PRL102 (2009) 242002]
 - $M=4143.4\pm 3.0\pm 0.6$ MeV
 - $\Gamma=15.3_{-6.1}^{+10.4} \pm 2.5$ MeV
 - Necessarily exotic since it is narrow and above the DsDs threshold
 - [$c s \bar{c} \bar{s}$] tetraquark ?
 - Hint of a second structure: $X(4274)$




$X(4140)$ and $X(4274)$



- Confirmed by CMS with large statistics
- But the background is also large



[PLB 734 (2014) 261]





Observation of four X




- With Run-1 $B^+ \rightarrow J/\psi\phi K^+$ data, LHCb performed 1st amplitude fit with 4300 signals
- Observed $X(4140)$, $X(4274)$, $X(4500)$ and $X(4700)$

[PRL118(2017)022003]




WEIGHTED AVERAGE
22+8-7 (Error scaled by 1.3)



The LHC as a Beauty and Charm factory

Proton-Proton Collisions at $\sqrt{s} = 13$ TeV
~ 20 000 $b\bar{b}$ pairs per second, x 20 of $c\bar{c}$ pairs




High B-baryon production fraction


$$\begin{array}{cccc} B^+ & : & B^0 & : & B_s^0 & : & \Lambda_b^0 \\ (u\bar{b}) & & (d\bar{b}) & & (s\bar{b}) & & (udb) \\ 4 & : & 4 & : & 1 & : & 2 \end{array}$$

LHCb detector and performance

JINST 3 (2008) S08005



- $2 < \eta < 5$ range: $\sim 25\%$ of $b\bar{b}$ pairs inside LHCb acceptance




[Int. J. Mod. Phys. A 30 (2015) 1530022]

Impact parameter:	$\sigma_{IP} = 20 \mu\text{m}$
Proper time:	$\sigma_\tau = 45 \text{ fs}$ for $B_s^0 \rightarrow J/\psi\phi$ or $D_s^+\pi^-$
Momentum:	$\Delta p/p = 0.4 \sim 0.6\%$ (5 – 100 GeV/c)
Mass :	$\sigma_m = 8 \text{ MeV}/c^2$ for $B \rightarrow J/\psi X$ (constrained $m_{J/\psi}$)
RICH $K - \pi$ separation:	$\epsilon(K \rightarrow K) \sim 95\%$ mis-ID $\epsilon(\pi \rightarrow K) \sim 5\%$
Muon ID:	$\epsilon(\mu \rightarrow \mu) \sim 97\%$ mis-ID $\epsilon(\pi \rightarrow \mu) \sim 1 - 3\%$
ECAL:	$\Delta E/E = 1 \oplus 10\%/\sqrt{E(\text{GeV})}$



LHCb collected luminosity

LHCb Cumulative Integrated Recorded Luminosity in pp, 2010-2018




Signal: Run2 = $4 \times$ Run1

We add Run2 data and improved selection for this analysis

B mass fit and background

[arXiv:2103.01803]

- 24k B^\pm signal and purity
96% in signal window
(± 15 MeV)
 - Signal: Hypatia function
 - Background: 2nd order polynomial



The largest $B^+ \rightarrow J/\psi\phi K^+$ sample so far

- In the B^\pm signal, about 2% are non- ϕ $B^\pm \rightarrow J/\psi K^+ K^- K^\pm$
 - They are neglected in the amplitude model but considered in the evaluation of the systematic uncertainties.



- Signal yield is **6×** larger than in the previous publication
- Background fraction **$\beta = 4\%$** is almost a factor of 6 smaller
- Signal efficiency is **15%** higher because of usage of PID in MVA



Dalitz plots

- In ± 15 MeV signal mass window
- Clearly visible: 4 structures in $J/\psi\phi$ mass and **an obvious $J/\psi K^+$ band**
- No clear $K^{*+} \rightarrow K^+\phi$ peaks because of K^{*+} resonances are broad?

[arXiv:2103.01803]





6D amplitude fit

[PRD95(2017)012002]




- Candidates in the signal region are used in the fit
- PDF includes **signal** and **background** components

$$\begin{aligned} -\ln L(\vec{\omega}) &= -\sum_i \ln [(1 - \beta) \mathcal{P}_{\text{sig}}(m_{\phi K i}, \Omega_i | \vec{\omega}) + \beta \mathcal{P}_{\text{bkg}}(m_{\phi K i}, \Omega_i)] \\ &= -\sum_i \ln \left[(1 - \beta) \frac{|\mathcal{M}(m_{\phi K i}, \Omega_i | \vec{\omega})|^2 \Phi(m_{\phi K i}) \epsilon(m_{\phi K i}, \Omega_i)}{I(\vec{\omega})} + \beta \frac{\mathcal{P}_{\text{bkg}}^u(m_{\phi K i}, \Omega_i)}{I_{\text{bkg}}} \right] \\ &= -\sum_i \ln \left[|\mathcal{M}(m_{\phi K i}, \Omega_i | \vec{\omega})|^2 + \frac{\beta I(\vec{\omega})}{(1 - \beta) I_{\text{bkg}}} \frac{\mathcal{P}_{\text{bkg}}^u(m_{\phi K i}, \Omega_i)}{\Phi(m_{\phi K i}) \epsilon(m_{\phi K i}, \Omega_i)} \right] + N \ln I(\vec{\omega}) . \end{aligned}$$

- Helicity formalism for full amplitude construction
- Each decay chain is described by 6 observables
 - Resonant mass, and 5 angles to better determine J^P
- Resonant lineshape: Breit-Wigner; simplified K-matrix or Flatté function for systematic studies

Start from run 1 model

- Run 1 model cannot fit well the data, due to increase of statistics
- Selection of resonance model is required






$K^{*+} \rightarrow \phi K^+$ model

9 $K(n^{2S+1}L_J)$ excitations

All $K(1^+)$	
2^1P_1	$K(1^+)$
2^3P_1	$K'(1^+)$
1^3P_1	$K_1(1400)$ Replace NR
All $K(2^-)$	
1^1D_2	$K_2(1770)$
1^3D_2	$K_2(1820)$
All $K(1^-)$	
1^3D_1	$K^*(1680)$
2^3S_1	$K^*(1410)$ Replace 3^3S_1
$K(2^+)$	
2^3P_2	$K_2^*(1980)$
$K(0^-)$	
2^1S_0	$K(1460)$

0^+ cannot decay to ϕK^+

- Based on Godfrey-Isgur model
- Compared to run-1 model
 - Add three below threshold resonances to replace two components
 - Other high mass states are not significant, used as systematic study (Extended model)








Test new exotics

- The K^* model still cannot well describe the data
- Test new exotic states (X and Z_{CS}^+) of different J^P
 - $1^+ Z_{CS}$ and $1^+ X$, giving the largest improvements, were first included.
 - In 2nd iteration, several states giving large fit improvements were included in the default model: a second Z_{CS} (either 1^+ or 1^-), 1^- and $2^- X$ states.
- The default model includes $9 K^* + 7 X + 1 X(NR) + 2 Z_{CS}$



Default model fit

- Data is well described by the model



Fit results

- New states: $Z_{cs}(4000)$, $X(4685) > 15\sigma$
 $Z_{cs}(4220)$, $X(4630) > 5\sigma$
 $X(4150) < 5\sigma$


[arXiv:2103.01803]

Fit fraction

Contribution	Significance [$\times\sigma$]	M_0 [MeV]	Γ_0 [MeV]	FF [%]
$X(2^-)$	Syst. included(Stat.)			
$X(4150)$	4.8 (8.7)	$4146 \pm 18 \pm 33$	$135 \pm 28^{+59}_{-30}$	$2.0 \pm 0.5^{+0.8}_{-1.0}$
$X(1^-)$				
$X(4630)$	5.5 (5.7)	$4626 \pm 16^{+18}_{-110}$	$174 \pm 27^{+134}_{-73}$	$2.6 \pm 0.5^{+2.9}_{-1.5}$
All $X(0^+)$				$20 \pm 5^{+14}_{-7}$
$X(4500)$	20 (20)	$4474 \pm 3 \pm 3$	$77 \pm 6^{+10}_{-8}$	$5.6 \pm 0.7^{+2.4}_{-0.6}$
$X(4700)$	17 (18)	$4694 \pm 4^{+16}_{-3}$	$87 \pm 8^{+16}_{-6}$	$8.9 \pm 1.2^{+4.9}_{-1.4}$
$NR_{J/\psi\phi}$	4.8 (5.7)			$28 \pm 8^{+19}_{-11}$
All $X(1^+)$				$26 \pm 3^{+8}_{-10}$
$X(4140)$	13 (16)	$4118 \pm 11^{+19}_{-36}$	$162 \pm 21^{+24}_{-49}$	$17 \pm 3^{+19}_{-6}$
$X(4274)$	18 (18)	$4294 \pm 4^{+3}_{-6}$	$53 \pm 5 \pm 5$	$2.8 \pm 0.5^{+0.8}_{-0.4}$
$X(4685)$	15 (15)	$4684 \pm 7^{+13}_{-16}$	$126 \pm 15^{+37}_{-41}$	$7.2 \pm 1.0^{+4.0}_{-2.0}$
All $Z_{cs}(1^+)$				$25 \pm 5^{+11}_{-12}$
$Z_{cs}(4000)$	15 (16)	$4003 \pm 6^{+4}_{-14}$	$131 \pm 15 \pm 26$	$9.4 \pm 2.1 \pm 3.4$
$Z_{cs}(4220)$	5.9 (8.4)	$4216 \pm 24^{+43}_{-30}$	$233 \pm 52^{+97}_{-73}$	$10 \pm 4^{+10}_{-7}$

Angular projections

Angles in $K^* \rightarrow \phi K$
decay chain are
described well by
the fit








$Z_{cs}(4000)^+$

[arXiv:2103.01803]

- Argand diagram gives further evidence of resonant character
 - Magnitude and phase evolved in the counter-clockwise direction

numbers are $m_{J/\psi K^+}$ in MeV

- $Z_{cs}(4000)^+$ can be clearly viewed in the two slices of $m_{J/\psi\phi}$






Comparison with BESIII

- BESIII experiment recently reported 5.3σ observation of a very narrow Z_{CS}^- in $D_S^- D^* + DD_S^{*-}$ mass distributions
- Their masses are close, but $Z_{CS}(4000)^+$ is $\sim 10\times$ broader
- Tests are applied:
 - Fix $Z_{CS}(4000)^+$ to BESIII's result; $2\ln L$ is worse by 160
 - Adding on top of the default model almost doesn't improve the fit likelihood
- No evidence that $Z_{CS}(4000)^+$ is the same as $Z_{CS}(3985)^-$ seen by BESIII

[arXiv:2011.07855]



$$m_{\text{pole}}(Z_{CS}(3985)^-) = (3982.5_{-2.6}^{+1.8} \pm 2.1) \text{ MeV}/c^2,$$
$$\Gamma_{\text{pole}}(Z_{CS}(3985)^-) = (12.8_{-4.4}^{+5.3} \pm 3.0) \text{ MeV}.$$



J^P analysis

- Rejection significance: $\sigma \sim \sqrt{\Delta(-2\ln L)}$, using $2\ln L$ difference between preferred and alternative hypothesis.
 - Previous observed four $X J^P$ are confirmed
 - $Z_{cs}(4000)$ and $X(4685)$ are 1^+
 - $X(4630)$ prefers 1^{-+} [exotic quantum number] over 2^{-+} by 3σ
 - $Z_{cs}(4220)$ can be 1^+ or 1^-

Systematic uncertainty included

J^P	0^+	0^-	1^+	1^-	2^+	2^-
X(4630)	6.7σ	5.3σ	5.8σ	prefer	5.9σ	3.0σ
X(4500)	prefer	18σ	18σ	18σ	18σ	18σ
X(4700)	prefer	18σ	18σ	18σ	14σ	17σ
X(4140)	14σ	15σ	prefer	14σ	13σ	14σ
X(4274)	18σ	18σ	prefer	18σ	18σ	18σ
X(4685)	16σ	16σ	prefer	15σ	16σ	15σ
Z_{cs}(4000)	-	17σ	prefer	17σ	15σ	16σ
Z_{cs}(4220)	-	8.6σ	prefer	2.4σ	4.9σ	5.7σ



J^{PC} of X states

- For $X \rightarrow J/\psi\phi$, S-wave decays: $J^{PC} = (0,1,2)^{++}$,
P-wave decays $J^{PC} = (0,1,2,3)^{-+}$
- We expect S-wave dominates and this is the case.
- **We are confident that our J^{PC} determination is right**
 - We have randomly assigned J^P for X states, and found the default results give the best fit
 - We can easily distinguish e.g. 0^{++} vs 1^{++} using the correlation between two decay angles of $J/\psi \rightarrow \mu^+\mu^-$ and $\phi \rightarrow K^+K^-$

0^{++} only has the following two terms, 1^{++} contains more terms

$$|H_0|^2$$

$$\sin^2 \theta_\ell \cos^2 \theta_h$$

$$|H_+|^2 + |H_-|^2$$




$$\frac{1}{4} (1 + \cos^2 \theta_\ell) \sin^2 \theta_h$$

X(4140)

- $X(4140) M = 4118 \pm 11_{-36}^{+19} \text{ MeV}$,
 $\Gamma = 162 \pm 21_{-49}^{+24} \text{ MeV}$

No evidence of a narrow threshold resonance at $J/\psi\phi$ in our data

- By comparing the samples, both structures are similar, but LHCb's $X(4140)$ peak height is lower than CMS (efficiency enhanced at threshold?)
- CMS should update their results on this channel with a (much) larger data sample, and more sophisticated analysis technique, than previously.








New X states are necessary

- Can improve angular distributions
- Comparing the Legendre angular moments of Run 1 model and updated model, new $X(4630)$ and $X(4685)$ are required

$$\langle P_\ell^U \rangle = \sum_{i=1}^{N_{\text{events}}} \frac{1}{\epsilon_i} P_\ell(\cos \theta)$$

<https://cds.cern.ch/record/2751229>



Background subtracted and efficiency corrected distribution





Systematic sources

- Many sources (see backup) are considered
- Here only discuss several important ones for modelling
 - Extended model including 5 more K^*
 - 1^+ vs $1^- Z_{CS}(4220)$
 - Additional X states with different J^P in the extended model, **no further X contribution $>5\sigma$**
 - NR shape, and additional 1^+ or 2^+ NR X contributions
 - Flatté function to parameterize $X(4140)$ or $Z_{CS}(4000)$ to replace BW function
 - Neglected no- ϕ contribution: 1) Change the ϕ mass window from $\pm 15\text{MeV}$ to $\pm 7\text{MeV}$, 2) sFit to subtract no- ϕ contribution is performed as alternative to cFit
 - Several K-Matrix models for K^*


Summary

- 4 new $J/\psi K^+$ and $J/\psi \phi$ structures observed in $B^+ \rightarrow J/\psi \phi K^+$ decays with 6 times data and much clean environment
 - **Two $Z_{cs}^+ \rightarrow J/\psi K^+$ with new quark contents $c\bar{c}u\bar{s}$ are observed**
 $1^+ Z_{cs}(4000)^+$, significance $> 15\sigma$ and a broad $Z_{cs}(4220)^+ > 5\sigma$
 - A new $1^+ X(4685)$ is $> 15\sigma$, and new $X(4630) > 5\sigma$
 - 4 X states previously observed are confirmed, and J^{PC} determined with higher significances
- Understanding of $Z_{cs}(4000)^+$ and $Z_{cs}(3985)^-$ may shed lights on molecular and compact tetraquarks





LHCb Upgrade I



CERN-LHCC-2011-001

Upgrade I: installation ongoing

- ❑ Almost a new detector for factor 5 luminosity increase
- ❑ Remove the **hardware trigger** → all detector read out at 40 MHz
- ❑ Expect to have data of **23 fb⁻¹** by 2024 and of **50 fb⁻¹** by 2029

- ❑ Efficiency of pure hadronic final states will be **3x** **7x** **Run1+2** **doubled**, good for studies of $(\eta_c, \chi_{cJ})(K, \phi)$ and $D_{(s)}^{(*)} \bar{D}_{(s)}^{(*)}$ to search for various J^P exotics

Thank you!



Expected yields in future

- We are now boosting our data to a new level
 - Expect to **7x** more data (**14x** more hadronic events) by 2029 than current data
 - Could have another factor of **6** increase from Upgrade II

Decay mode	LHCb		
	23 fb ⁻¹	50 fb ⁻¹	300 fb ⁻¹
$B^+ \rightarrow X(3872)(\rightarrow J/\psi \pi^+ \pi^-) K^+$	14k	30k	180k
$B^+ \rightarrow X(3872)(\rightarrow \psi(2S)\gamma) K^+$	500	1k	7k
$B^0 \rightarrow \psi(2S) K^- \pi^+$	340k	700k	4M
$B_c^+ \rightarrow D_s^+ D^0 \bar{D}^0$	10	20	100
$\Lambda_b^0 \rightarrow J/\psi p K^-$ [*]	680k	1.4M	8M
$\Xi_b^- \rightarrow J/\psi \Lambda K^-$	4k	10k	55k
$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$	7k	15k	90k
$\Xi_{bc}^+ \rightarrow J/\psi \Xi_c^+$	50	100	600


[*] updated according to the latest result

CERN-LHCC-2018-027
arXiv:1808.08865



Significance

- Use $\text{ndf} = 2 \times N$ of parameters for new resonance
- Verified by pseudo-experiments



500 toy samples without $Z_{CS}(4000)$ are generated.

The significance obtained from the tail extrapolation to the data is 15.2σ , which is consistent with 15.7σ obtained from the empirical method using the χ^2 PDF with ndf equal to twice the number of additional free parameters

$$n_\sigma = \sqrt{2} \text{TMath::ErfcInverse} \left(f \cdot \frac{\text{TMath::Prob}(\Delta_{2\ln\mathcal{L}}^{\text{data}}, \text{NDF})}{\text{TMath::Prob}(\Delta_{2\ln\mathcal{L}}^0, \text{NDF})} \right)$$




Run-1 results

Contribution	Significance	Fit results		
		M_0 [MeV]	Γ_0 [MeV]	FF%
All K(1 ⁺)	8.0 σ			$42 \pm 8_{-9}^{+5}$
NR _{ϕK}				$16 \pm 13_{-6}^{+35}$
2 ¹ P ₁ K(1 ⁺)	7.6 σ	$1793 \pm 59_{-101}^{+153}$	$365 \pm 157_{-215}^{+138}$	$12 \pm 10_{-6}^{+17}$
2 ³ P ₁ K'(1 ⁺)	1.9 σ	$1968 \pm 65_{-172}^{+70}$	$396 \pm 170_{-178}^{+174}$	$23 \pm 20_{-29}^{+31}$
All K(2 ⁻)	5.6 σ			$11 \pm 3_{-5}^{+2}$
1 ¹ D ₂ K ₂ (1770)	5.0 σ	$1777 \pm 35_{-77}^{+122}$	$217 \pm 116_{-154}^{+221}$	
1 ³ D ₂ K ₂ (1820)	3.0 σ	$1853 \pm 27_{-35}^{+18}$	$167 \pm 58_{-72}^{+83}$	
1 ³ D ₁ K(1 ⁻)				
K*(1680)	8.5 σ	$1722 \pm 20_{-109}^{+33}$	$354 \pm 75_{-181}^{+140}$	$6.7 \pm 1.9_{-3.9}^{+3.2}$
2 ³ P ₂ K(2 ⁺)				
K*(1980)	5.4 σ	$2073 \pm 94_{-240}^{+245}$	$678 \pm 311_{-559}^{+1153}$	$2.9 \pm 0.8_{-0.7}^{+1.7}$
3 ¹ S ₀ K(0 ⁻)				
K(1830)	3.5 σ	$1874 \pm 43_{-115}^{+59}$	$168 \pm 90_{-104}^{+280}$	$2.6 \pm 1.1_{-1.8}^{+2.3}$
All X(1 ⁺)				$16 \pm 3_{-2}^{+6}$
X(4140)	8.4 σ	$4146.5 \pm 4.5_{-2.8}^{+4.6}$	$83 \pm 21_{-14}^{+21}$	$13.0 \pm 3.2_{-2.0}^{+4.8}$
X(4274)	6.0 σ	$4273.3 \pm 8.3_{-3.6}^{+17.2}$	$56 \pm 11_{-11}^{+8}$	$7.1 \pm 2.5_{-2.4}^{+3.5}$
All X(0 ⁺)				$28 \pm 5 \pm 7$
NR _{J/$\psi$$\phi$}	6.4 σ			$46 \pm 11_{-21}^{+11}$
X(4500)	6.1 σ	$4506 \pm 11_{-15}^{+12}$	$92 \pm 21_{-20}^{+21}$	$6.6 \pm 2.4_{-2.3}^{+3.5}$
X(4700)	5.6 σ	$4704 \pm 10_{-24}^{+14}$	$120 \pm 31_{-33}^{+42}$	$12 \pm 5_{-5}^{+9}$



LHCb Upgrade II



Upgrade II: started to investigate


- ❑ Aim to collect $> 300 \text{ fb}^{-1}$
- ❑ Instantaneous $\mathcal{L} = 2 \times 10^{34}$, x10 with respect to Upgrade I
- ❑ Expression of Interest issued in 2017 [[CERN-LHCC-2017-003](#)]
- ❑ Physics case document released [[CERN-LHCC-2018-027](#)]
- ❑ Green light from LHCC to proceed to TDRs (expected ~late 2020)



Detector performance

Vertexing


B_s^0 oscillations with $B_s^0 \rightarrow D_s \pi$



[New J. Phys. 15 (2013) 053021] [EPJ C73 (2013) 2431]


PID

K/π ID efficiency and misID rate



Tracking

$\mu^+ \mu^-$ mass spectrum



[PRL 111 (2013) 101805]

Impact parameter:

Proper time:

Momentum:

Mass:

RICH $K - \pi$ separation:

Muon ID:

ECAL:

$$\sigma_{IP} = 20 \mu\text{m}$$

$$\sigma_{\tau} = 45 \text{ fs for } B_s^0 \rightarrow J/\psi\phi \text{ or } D_s^+ \pi^-$$

$$\Delta p/p = 0.4 \sim 0.6\% (5 - 100 \text{ GeV}/c)$$

$$\sigma_m = 8 \text{ MeV}/c^2 \text{ for } B \rightarrow J/\psi X \text{ (constrained } m_{J/\psi})$$

$$\epsilon(K \rightarrow K) \sim 95\% \text{ mis-ID } \epsilon(\pi \rightarrow K) \sim 5\%$$

$$\epsilon(\mu \rightarrow \mu) \sim 97\% \text{ mis-ID } \epsilon(\pi \rightarrow \mu) \sim 1 - 3\%$$

$$\Delta E/E = 1 \oplus 10\%/\sqrt{E(\text{GeV})}$$



Systematic uncertainty

- To evaluate uncertainties due to the **fixed masses and widths of known K^*** resonances: free the masses and widths but impose Gaussian constraints to the PDG values.
- χ_{IP}^2 of B^+ is not well modeled, smeared to match the data.
- To explore uncertainty in the background model, vary the B^+ sideband window.
- The uncertainty in the background fraction β : change background shape to exponential function.
- Vary the Blatt-Weisskopf barrier factor d (hadron-size parameter).
- **Vary the smallest allowed orbital momentum** in the resonance description function, associate the L dependent term with each LS coupling.

$$R_{K_n^*}(m_{K\phi}) = \underbrace{B'_{L_B^{K_n^*}}(p, p_0, d) \left(\frac{p}{M_B}\right)^{L_B^{K_n^*}}}_{\text{Angular momentum barrier factor}} \underbrace{\text{BW}(m_{K\phi} | M_0^{K_n^*}, \Gamma_0^{K_n^*})}_{\text{Relative Breit-Wigner function}} \underbrace{B'_{L_{K_n^*}}(q, q_0, d) \left(\frac{q}{M_0^{K_n^*}}\right)^{L_{K_n^*}}}_{\text{Orbital momentum}}$$



Systematic uncertainty

- Uncertainty due to the choice of NR component, change the constant parameterization to exponential function.
- 1^+ or 2^+ NR X contributions are optionally introduced.
- The difference between nominal model and extended model.
- Flatté function to parameterize $X(4140)$ or $Z_{CS}(4000)$ to replace BW function.

$$\text{Flatte}_X(m|M_0, g_{J/\psi\phi}, g_{D_s^*D_s}) = \frac{1}{M_0^2 - m^2 - iM_0(g_{J/\psi\phi}\rho_{J/\psi\phi} + g_{D_s^*D_s}\rho_{D_s^*D_s})},$$

- Additional X states with different J^P in the extended model.
- Neglected no- ϕ contribution: 1) Change the ϕ mass window from $\pm 15\text{MeV}$ to $\pm 7\text{MeV}$, 2) sFit to subtract no- ϕ contribution is performed as alternative to cFit
- Modification of K^* width: as the partial width to ϕK is unknown, try a fit with mass dependence of the width driven by the lowest allowed decay channel, which is $K\pi$ for the natural spin-parity and $K\omega$ for others.



Systematic uncertainty

- As an alternative to the 2D factorization of 6D background PDF, decompose the background density into multidimensional moments in the K^* decay chain variables (this uncertainty is small)
- K-Matrix model :
 1. Some K^* with the same J^P are overlapping, we use a simple K-Matrix formula to describe them as alternative

$$RKM_n(m|M_{0n}, \Gamma_{0n}) = \frac{1}{1 - i \left(\sum_j \frac{M_{0j} \Gamma_{0j}(m)}{M_{0j}^2 - m^2} + f_{sc} \cdot \rho(m) \right)},$$

denominator sums over the same J^P K^* resonances, f_{sc} accounts for possible non-resonance contribution. This fit didn't change the conclusion.

2. Alternative K-Matrix model with two coupling channels are tested, used to describe the $2^1 P_1$ and $2^3 P_1$ K^* resonances

$$\mathcal{K}_{ba}(s) = \sum_R \frac{g_b^R g_a^R}{M_R^2 - s} + \sum_{i=0}^{N_{b.g.}} b_{ba}^{(i)} s^i.$$

more floating parameters are included, the nominal model is stable.



Table 2: Summary of the systematic errors on the parameters of the $Z_{cs}(4000)^+$ and $X(4685)$ states. All numbers for masses and widths are in MeV and fit fractions in %.

Source	$Z(4000)$			$X(4685)$		
	M_0	Γ_0	FF	M_0	Γ_0	FF
Fixed M_0 & Γ_0	-0.22	-3.60	-0.83	-0.14	2.72	0.25
χ^2_{IP} smearing	0.21	1.01	0.09	-0.53	1.11	0.12
Right sideband	0.01	0.58	0.11	-0.13	1.07	-0.13
Left sideband	-0.30	-1.16	-0.24	-0.09	-2.21	0.09
$\beta = 0.043$	-0.06	-0.00	0.01	0.01	-0.70	-0.09
$\beta = 0.037$	-0.02	0.26	0.02	-0.33	0.21	0.03
L0 Trigger	0.45	0.58	0.19	-0.58	1.12	0.11
PID efficiency	-1.06	-1.82	-0.69	-0.82	-4.42	-0.26
MC size	2.39	9.93	1.54	3.02	7.00	0.65
ϕ window	-4.71	-23.91	-2.75	8.60	-26.60	-1.17
Non ϕ subtraction	-2.87	-18.39	-1.79	12.40	-39.80	-1.80
Poly NR	-4.24	-16.36	-2.56	4.26	-22.07	-1.28
X NR(1^+)	1.49	-21.25	-2.53	-15.72	35.54	3.84
X NR(2^+)	2.16	3.09	1.26	1.88	-6.87	-0.03
BW $d=1.5$	-0.29	-5.27	-0.58	0.29	1.55	2.14
BW $d=4.5$	0.08	1.81	0.04	0.06	-3.53	-1.06
L	2.75	-3.19	-1.18	2.45	-24.33	-1.48
$X(4140)$ Flatté	0.52	-2.80	-0.45	-3.77	15.14	1.37
Extended model	-2.35	-6.66	-1.16	-3.61	-6.53	-0.94
Additional X	-0.68	2.07	0.30	0.74	-3.11	-0.18
$1^- Z$	-14.00	-21.09	-3.46	-9.41	-5.60	-1.52
K^* BW	0.08	-0.66	-0.32	-0.06	-8.09	-0.82
K-Matrix	-3.75	-20.80	-2.85	4.10	-11.95	-0.06
$Z_{cs}(4000)$ Flatté	0.18		2.83	-0.85	2.79	0.18
Background model	0.10	-0.32	-0.12	-1.04	-1.72	-0.15
Total	(-14.26 , +3.85)	(-26.26 , +26.26)	(-3.43 , +3.41)	(-16.05 , +12.82)	(-40.85 , +36.72)	(-1.96 , +3.92)

Thresholds vs LHCb run1 data

arXiv:2101.01021

