



Introductory Part

Collider Physics

Toni Baroncelli
Haiping Peng
USTC

Year 2023



Practicalities

These slides (and the Lecture Notes) will be made available at an Indico page being prepared

<http://cicpi.ustc.edu.cn/indico/categoryDisplay.py?categId=309>

Attending a course in English is difficult for young persons who are not too familiar with foreign languages

*I understand your difficulty and appreciate your effort
The world of HEP (High Energy Physics) is a world-wide collaboration and English is the
standard tool of communication. Attending these lectures will help you to improve your foreign
language skills*

There are several ways to contact me:

1. By sending a mail to toni.baroncelli@cern.ch, please indicate “Student of Collider Physics” in the subject
2. Knock at my Office, 6th floor of the Modern Physics building (A606)

Do not hesitate to contact me for any question concerning the course

At the beginning of each new lecture I will give a short summary of the previous lecture with a list of main physics points

Lecture notes will be prepared for (most) of the course




Lecture Notes



Collider Physics

Haiping Peng
Toni Baroncelli



Lecture Notes for the 2022
"Collider Physics"
Course at USTC

I have prepared Lecture Notes for most of the course
(.. Dynamic document, it will evolve with passing lectures).


Freely available



Collider Physics Course, Year 2021

- The points on the plot are seen to group in two families, corresponding to "Lepton Colliders" and "Hadron Colliders". Both families exhibit a similar growth as a function of time.
- "Lepton Colliders" (in practice today we only accelerate electrons and positrons), however, are significantly below the "Hadron Colliders" family. This is due to difficulties in the acceleration of electrons that limit the maximum centre of mass energy achievable.
- The linear trend observed in years 1960 to 2000 seems to curb due to technological limits both for hadron and lepton machines.
- The Livingston plot is also populated by 'future' accelerators indicating that, even though slower, the evolution of accelerators is expected to make available new generations of machines within a timescale of ~thirty years.
- All this indicates that it will be more and more difficult to access new accelerators and that once a new machine is built this will have to be exploited for decades.

8.



9. Figure 2-3: The Livingston plot.

As we have seen, in the last half a century, the physicists have been able to gain a factor 10 in \sqrt{s} by colliding two head-on beams. The complex technology that is at the base of these very accurate probes is perhaps at the limit of its reach. Also, the cost of these colliders is exploding at the level that no

Country alone in the world can sustain the cost of an accelerator of the future.


10. We all hope we will continue to have access to more and more powerful machines, but at some point, we will need a new technology breakthrough, since not many funds will be easily available in the foreseeable future.



Collider Physics Course, Year 2021

term $E_M^2 (1 - \cos\theta)$ puts in correlation the scattering angle and the scattering energy of

the electron. If we increase the target mass M the energy transfer $E-E'$ increases as well.




32. Figure 7-3 Ratio E'/E for two different values of the electron energy (0.5 and 10 GeV) and for two values of the atomic number

The correlation plot between θ and the ratio E'/E is shown in the Figure 7-3 for two different values of the electron energy (0.5 and 10 GeV) and for two values of the atomic

number ($A=1$, hydrogen, $A=50$, tin). While for hydrogen the ratio E'/E is seen to vary very significantly, going from 0.6 to less than 0.1 at large angles, for high mass values the variation is much less important.

7.2. Rutherford scattering

First scattering experiments tried to give an answer to a basic (for the time) question: is the target of our scattering experiment a point-like object? What is the size of the scattering centres of our experiment? The approach followed by Rutherford was that of taking a source of α particles (easy, use a radio-active source) and collimated emitted α particles against a thin foil of Gold (Au). Later physicists realised that the results of the Rutherford experiment were interesting but could not be used beyond some level: using




33. Figure 7-4 kinematics of the α -p scattering

electrons (a real point-like projectile) was much better and the results could be used to study structures much smaller than with complex and massive α particles.

The kinematics of the scattering process is sketched in Figure 7-4. The projectile travels at a transverse distance b (see Figure 7-5)



Collider Physics Course, Year 2021: Deep inelastic scattering



62. Figure 9-9 Evolution of F_2 as a function of Q^2

10.3. Measuring (= Counting) the number of colours⁷

When the concept of colour was introduced above, we said that there are three colours, red, green and blue. This is not an information that comes down from the sky, it descends from an experimental observation that is described below.

This experimental observation is based on




Figure 8

the study the production of

- $q\bar{q}$ pairs and of
- $\mu^+\mu^-$ pairs

in e^+e^- interactions where only virtual photons can be exchanged during the interaction between electrons and positrons.

The production of $q\bar{q}$ pairs is shown in top part of figure 8. In the bottom part the Feynman diagrams for a $\mu^+\mu^-$ pairs creation is shown (mostly mediated by photons, the exchange of the Z boson contributing very little).

In both final states, $q\bar{q}$ pairs and $\mu^+\mu^-$ pairs, the cross section depends on α_{em}^2 . Let us write the formula for the cross section in both cases:

$$\sigma(e^+e^- \rightarrow \mu^+\mu^-) = \frac{4\pi\alpha_{em}^2\hbar^2}{3s}$$

In this expression the charge squared of the muon, q_μ^2 , has been omitted since it is equal to 1. Things are different in the hadronic final states, which is the superposition of many different states,

$$\sigma(e^+e^- \rightarrow \gamma \rightarrow q\bar{q}) = N_c \frac{4\pi\alpha_{em}^2\hbar^2}{3s} \sum_{q=1}^{N_f} q_q^2$$

For both the final states the structure is the same but in the hadronic many more possible final states are open: this number is just given by $N_c \sum_{q=1}^{N_f} q_q^2$ where N_c is the number

⁷ Section 9.2 of [3]



Overview of the Course

The Course will be shared between

- Toni Baroncelli (1st part → 2 months)
- And Haiping Peng (2nd part → 2 months)


And is organized in several parts, lasting in total 16 weeks + 2 weeks for examinations.

Topic	Weeks	Who	from	→	# lectures
Introduction to basic concepts	2	T.Baroncelli	08/03/23	17/03/23	4
Deep Inelastic Scattering	1	T.Baroncelli	22/03/23	24/03/23	6
Accelerators	1	T.Baroncelli	29/03/23	31/03/23	8
Detectors	1	T.Baroncelli	05/04/23	07/04/23	10
Measurements at Colliders	3	T.Baroncelli	12/04/23	28/04/23	16
Standard Model Theory	2	H.Peng	03/05/23	12/05/23	4
CPV theory and experiment (BELLE, BABAR, LHCb)	2	H.Peng	17/05/23	26/05/23	8
Hadron physics (BESIII, STCF)	2	H.Peng	31/05/23	09/06/23	12
Higher Symmetries (GUT, SUSY, Superstrings....)	2	H.Peng	14/06/23	23/06/23	16
	16				



Forward: (Particle) Physics?

Experimental Science! Observation of *different* variables $1 \dots n+3$



Theory: establishes relations among observables (experiments)

- All observables are accounted for in the theory \rightarrow numerical predictions;
- Errors in the measurement;
- Infinite theories may describe the same set of observables;
- Two different theories are different IF they predict different observables or different outcomes of the same observables;
- \rightarrow consistency of the theory
- \rightarrow accuracy in the prediction;
- A theory can be falsified because of wrong predictions (... better say 'incompatible with observables');
- A theory cannot be qualified (proven) because a newer experiment can always disprove an older theory \rightarrow THEORIES EVOLVE !




Forward: why 'Collider Physics'?

We will see that to access the intimate structure of matter we have to use probes with wave-lengths as small as possible → high energy!

1. Accelerate particles onto targets (used in the past) or
2. Collide two beams against each other

The second option became accessible only with ~modern technologies.


Unprecedented high energies are reached → many discoveries



Maximilien Brice (CERN) - CERN Document Server, CC BY-SA 3.0, <https://commons.wikimedia.org/w/index.php?curid=29027732>



LHC tunnel



CMS Experiment



Part One - Structure

Topics

1. Probing the Structure of Matter
 2. Constituents of Matter & Quantum Numbers
 3. The Standard Model, Interactions and Vector Bosons
 4. Symmetries and Conservation Laws
 5. The Electro-Magnetic Case
 6. Feynman diagrams
 7. Cross Sections & the Golden Rule
 8. Electron – Nucleus Scattering
 9. Rutherford Scattering
 10. Form Factors
- The way we see things today,
no historical approach → SM
- Basic Ingredients
- First Experiments



Prologue: Many Order of Magnitude

(Reduced) Planck's Constant ($\hbar = h/2\pi$) h

The uncertainty principle:
 "position x (with uncertainty Δx) and momentum p_x (with uncertainty Δp_x) cannot simultaneously be known to better than

$$\Delta x \Delta p_x \sim \hbar/2.$$


A relation for the energy is obtained by multiplying c ,

$$\Delta x \Delta E \sim \frac{\hbar c}{2}$$

which gives numerically,

$$\Delta E (MeV) = \frac{1.973^{-11} (MeV \text{ cm})}{2 \Delta x (cm)}$$

Also $\Delta x = c \Delta t \rightarrow \Delta t \Delta E \sim \frac{\hbar}{2}$





Prologue: the Quest for High Energy

- Discovery range is limited by available data, i.e. by resources (like a microscope).
- The true variable is the resolving power of our microscope.

Quantum Mechanics, wave-particle duality:

- a particle has wave-like properties. A wave is characterized by its wavelength;
- a wave has particle-like properties.; a particle is characterized by its energy or its momentum.

The larger the particle energy, the smaller the associated wavelength.




Prologue: the Quest for High Energy

- Resolving Power \propto Energy transferred in interaction, $1/\sqrt{Q^2}$ [i.e. $\propto 1/\sqrt{s}$, the CM energy]
- For *non point-like objects*, replace \sqrt{s} with the CM energy at component level, $\sqrt{\hat{s}}$ ($\sqrt{\hat{s}} < \sqrt{s}$) (quarks in a proton, will see later).

Δx (cm)	E	Tool
10^{-5}	2 eV	Microscopes
10^{-8}	2 keV	X rays
10^{-11}	2 MeV $\simeq 4m_e$	γ rays
10^{-14}	2 GeV $\simeq 2m_p$	Accelerators
10^{-16}	200 GeV $\simeq 2m_{W,Z}$	Accelerators
10^{-17}	2 TeV	Accelerators

LHC, as an example




The CM energy at component level is lower! Quarks carry a fraction of the proton energy



Prologue: the Quest for High Energy

- In the last half a century, the physicists have been able to gain a factor 10 in \sqrt{s} every 10 years (see the "Livingston plot").
 - Fixed target (1 beam on a thin target)
 - Two head-on beams (more complex technology)
- **No Country** in the world can sustain the cost of an accelerator of the future
- Hope it will continue like that, but needs IDEAS, since not many \$\$\$ (or €€€) will be available.


CEPC in China(*)?



(*) <http://cepc.ihep.ac.cn/intro.html>




The Livingston Plot





Prologue: The Standard Model

- SM designates the theory of the Electromagnetic, Weak and Strong interactions. The theory has grown in time, the name went together.
- The development of the SM is an interplay between new ideas and measurements.



- Many theoreticians contributed : since the G-S-W (S.Glashow, A.Salam,S.Weinberg) model is at the core of the SM, it is common to quote them as the main authors.



Reminder (mostly) of Quantum Mechanics

Look at Lecture Notes!

- Contravariant four vectors (index up) $x^\mu = (t, x, y, z)$
- Covariant four vectors (index down) $x_\mu = (t, -x, -y, -z)$
- $x^\mu x_\mu = x^0 x_0 + x^1 x_1 + x^2 x_2 + x^3 x_3$

QM = Quantum Mechanics
 \hbar = Planck Constant

- all the information regarding a physical system is contained in the corresponding wavefunction
- Free particles (we use natural units $\hbar, c = 1$) can be expressed as

$$\psi(\mathbf{x}, t) = N \cdot e^{i(\mathbf{p} \cdot \mathbf{x} - Et)}$$

- Operators acting on a wavefunction return an observable (real number) $\hat{A}\psi = a\psi$
- Hamiltonian: $E = H = T + V$

Natural Units

Quantity	kg, m, s	\hbar, c, GeV	$\hbar = c = 1$
Energy	$\text{kg m}^2 \text{s}^{-2}$	GeV	GeV
Momentum	kg m s^{-1}	GeV/c	GeV
Mass	kg	GeV/c^2	GeV
Time	s	$(\text{GeV}/\hbar)^{-1}$	GeV^{-1}
Length	m	$(\text{GeV}/\hbar c)^{-1}$	GeV^{-1}
Area	m^2	$(\text{GeV}/\hbar c)^{-2}$	GeV^{-2}

- If an Hamiltonian leaves unchanged an observable then $[\hat{H}, \hat{O}] = \hat{H}\hat{O} - \hat{O}\hat{H} = 0$
- $\hat{p} = -i\nabla$ and $\hat{E} = i\frac{\partial}{\partial t}$
- \hat{p} returns \mathbf{p} , \hat{E} returns E



Non-relativistic Schrödinger equation

$$E = H = T + V = \frac{\mathbf{p}^2}{2m} + V \sim \text{Valid at low energy}$$

$\hat{p} = -i\nabla$ and $\hat{E} = i\frac{\partial}{\partial t}$ applied to $\psi(\mathbf{x}, t) = N \cdot e^{i(\mathbf{p}\cdot\mathbf{x} - Et)}$ gives

$$\rightarrow \hat{H}\psi(\mathbf{x}, t) = \frac{1}{2m} (-i\nabla)^2 \psi(\mathbf{x}, t) + \hat{V}\psi(\mathbf{x}, t)$$

$$i \frac{\partial \psi(\mathbf{x}, t)}{\partial t} = -\frac{1}{2m} \frac{\partial^2 \psi(\mathbf{x}, t)}{\partial x^2} + \hat{V}\psi(\mathbf{x}, t)$$

Two 'weak' points:

- it contains 2nd order derivatives in space and 1st order in time → it is not relativistic invariant → inadequate to describe high energy phenomena (low energy approximation)
- Has some other theoretical weakness



The Klein-Gordon Equation

The *Schrödinger* equation is not relativistic invariant → go to the Einstein relation between energy and momentum:

$$E = \frac{\mathbf{p}^2}{2m} \rightarrow E^2 = \mathbf{p}^2 + m^2$$

(Note that you go from E to E^2)

The E^2 and \mathbf{p}^2 terms of this equation are interpreted as operators that act on a wavefunction (as before $\hat{p} = -i\nabla$ and $\hat{E} = i \frac{\partial}{\partial t}$):

$$\hat{E}^2 \psi(x) = \hat{\mathbf{p}}^2 \psi(x) + m^2 \psi(x)$$

$$(\partial^\mu \partial_\mu + m^2) \psi(x, t) = 0$$

Where

$$\partial^\mu \partial_\mu = \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2}$$

Problem is that the solution gives $E = \pm \sqrt{p^2 + m^2}$
(and other theoretical difficulties)

It gives an unphysical negative energy solution !

Alternative solution needed



The Dirac Equation

Dirac tried a different approach and wrote an equation with first order derivatives only

$$\hat{E}\psi(\mathbf{x}, t) = (\boldsymbol{\alpha} \cdot \hat{\mathbf{p}} + \beta \cdot m) \cdot \psi(\mathbf{x}, t)$$

Attention!! Terms α and β are NOT necessarily numbers!

If we write this explicitly (using the usual replacement $\hat{p} = -i\nabla$ and $\hat{E} = i\frac{\partial}{\partial t}$) we get

$$i\frac{\partial}{\partial t}\psi = \left(-i\alpha_x\frac{\partial}{\partial x} - i\alpha_y\frac{\partial}{\partial y} - i\alpha_z\frac{\partial}{\partial z} + \beta m\right)\psi$$

$$E^2 = \mathbf{p}^2 + m^2$$

The solutions of the Dirac equation have to be a solution of the Klein-Gordon equation. → Constraints on the α and β terms.

$$\begin{aligned} \alpha_x^2 &= \alpha_y^2 = \alpha_z^2 = I \\ \alpha_j\beta + \beta\alpha_j &= 0 \\ \alpha_j\alpha_k + \alpha_k\alpha_j &= 0 \quad (j \neq k) \end{aligned}$$

Where I represents unity. → α and β are matrices. It can be shown that α and β matrices, have even dimensions, the minimum value being 4x4

- have trace 0
- have eigenvalues ± 1
- are Hermitian

the solution for the Dirac equation has to be a four-component (*Dirac-spinor*) wavefunction:

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix}$$



Physical Interpretation – Association to Spin

It is natural to think of spinors as associated to spin

$$u_1(E, \mathbf{p}) = N \begin{pmatrix} 1 \\ 0 \\ \frac{p_z}{E+m} \\ \frac{p_x+ip_y}{E+m} \end{pmatrix} \text{ and } u_2(E, \mathbf{p}) = N \begin{pmatrix} 0 \\ 1 \\ \frac{p_x-ip_y}{E+m} \\ \frac{-p_z}{E+m} \end{pmatrix}$$


Positive energy solution, u_1 and u_2 wave functions with positive energy and spin up and spin down respectively

$$u_3(E, \mathbf{p}) = N \begin{pmatrix} \frac{p_z}{E-m} \\ \frac{p_x+ip_y}{E-m} \\ 1 \\ 0 \end{pmatrix} \text{ and } u_4(E, \mathbf{p}) = N \begin{pmatrix} \frac{p_x-ip_y}{E-m} \\ \frac{-p_z}{E-m} \\ 0 \\ 1 \end{pmatrix}$$

Negative energy solution, u_3 and u_4 wave functions with *negative* energy and spin up and spin down respectively

These solutions satisfy both the Einstein relation $E^2 = p^2 + m^2$ and the Dirac equation.

- The first two solutions have $E = \sqrt{p^2 + m^2}$ while
- the other two solutions have $E = -\sqrt{p^2 + m^2}$.



Again negative energy solutions !!




Physical Interpretation – Negative Energies?

The existence of negative energy solutions is a problem.

Dirac tried to solve it by saying that all negative energy states are ‘occupied’ and there is no room to accept ‘positive states’. → Historical interest only

Later *Stückelberg* and *Feynman* proposed a different interpretation:



The time dependence of the solutions to the Dirac equation is


$$e^{-iEt}$$

term. If you change

$$t \rightarrow -t \text{ and } E \rightarrow -E$$

the time behaviour is unaffected.

Solution → existence of an antiparticle with positive energy, opposite charge to that of the corresponding particle and propagating forward in time.



This interpretation was later validated by many experimental observations.



Spin and Helicity

At this point we may introduce two antiparticle spinors $v_1(E, \mathbf{p})$ and $v_2(E, \mathbf{p})$ where, simply,
the sign of E and \mathbf{p} have been reversed:


$$v_1(E, \mathbf{p})e^{-i(\mathbf{p}\cdot\mathbf{x}-Et)} = u_4(-E, -\mathbf{p})e^{-i(-\mathbf{p}\cdot\mathbf{x}-(-E)t)}$$

$$v_2(E, \mathbf{p})e^{-i(\mathbf{p}\cdot\mathbf{x}-Et)} = u_3(-E, -\mathbf{p})e^{-i(-\mathbf{p}\cdot\mathbf{x}-(-E)t)}$$

If we choose $p_z \neq 0., p_{x,y} = 0.,$ (as an example!) the spinors for particles and anti particles become

$$u_1(E, \mathbf{p}) = N \begin{pmatrix} 1 \\ 0 \\ \frac{p_z}{E+m} \\ 0 \end{pmatrix} \text{ and } u_2(E, \mathbf{p}) = N \begin{pmatrix} 0 \\ 1 \\ 0 \\ \frac{-p_z}{E+m} \end{pmatrix}$$

$$v_1(E, \mathbf{p}) = N \begin{pmatrix} \frac{p_z}{E-m} \\ 0 \\ 1 \\ 0 \end{pmatrix} \text{ and } v_2(E, \mathbf{p}) = N \begin{pmatrix} 0 \\ \frac{-p_z}{E-m} \\ 0 \\ 1 \end{pmatrix}$$





Spin and Helicity

If we define a spin-operator and we apply it to the spinors just introduced we get

$$\hat{S}_z = 1/2 \begin{pmatrix} \sigma_z & 0 \\ 0 & \sigma_z \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$\hat{S}_z u_1(E, 0.0, \pm p) = +1/2 u_1(E, 0.0, \pm p) \\ \hat{S}_z u_2(E, 0.0, \pm p) = -1/2 u_2(E, 0.0, \pm p) \\ \hat{S}_z v_1(E, 0.0, \pm p) = +1/2 v_1(E, 0.0, \pm p) \\ \hat{S}_z v_2(E, 0.0, \pm p) = -1/2 v_2(E, 0.0, \pm p)$$

A generalisation has to be introduced by defining the concept of helicity h :

$$h = \frac{\mathbf{S} \cdot \mathbf{p}}{p}$$

The helicity h is the component of the spin along the direction of motion. The corresponding operator \hat{h} is now defined as (see the 'Matrix section' in the Lecture Notes)

$$\hat{h} = 1/2 \begin{pmatrix} \sigma \cdot \hat{p} & 0 \\ 0 & \sigma \cdot \hat{p} \end{pmatrix}$$

- The helicity commutes with the free-particle Hamiltonian \rightarrow the spinors are both eigenstates of the helicity operator and of the free particle Hamiltonian.
- The corresponding eigenvalues are $\pm 1/2$.
- Particles with helicity $+1/2$ are righthanded, particles with helicity $-1/2$ are lefthanded.




The Standard Model

The Standard Model (SM) is the best description we have today of the microscopic world:

- Describes accurately known phenomena
- Important predictive power
- Incorporates known particles, forces and the interaction among them
- Predicted the existence of the Higgs Boson (discovered in 2012).

However the SM is not really a theory, it is rather a 'Model':

- Many parameters(*) have to be fixed 'by-hand' to describe data
- *SM* created an infrastructure with locations for particles and forces but is not able to explain why it is like that



The *SM* is not the end of the story!
More has to exist and needs to be discovered.

→ *BSM* is called the '*Beyond Standard Model*' Physics. It will incorporate the *SM* and its capacity to describe / predict microscopic phenomena

(*) SM parameters: masses of twelve fermions, three strengths of gauge interactions, two parameters for the Higgs potential, eight parameters of the mixing matrix CKM → 25 parameters



Constituents of Matter (a 'picture of the World')

Table 1 : Fundamental fermions and bosons in the standard model of the microcosm

Fermions

$\begin{pmatrix} u \\ d \end{pmatrix}$	$\begin{pmatrix} c \\ s \end{pmatrix}$	$\begin{pmatrix} t \\ b \end{pmatrix}$	Quarks
$\begin{pmatrix} \nu_e \\ e^- \end{pmatrix}$	$\begin{pmatrix} \nu_\mu \\ \mu^- \end{pmatrix}$	$\begin{pmatrix} \nu_\tau \\ \tau^- \end{pmatrix}$	Leptons

First family Second family Third family

Leptons and quarks considered to be the *ultimate* fermionic constituents of matter. Quarks (and antiquarks) appear in three different colours (and anticolours)

Bosons

Fundamental Interactions Mediators

Strong
Electromagnetic
Weak

8 gluons
 γ
 W^+, W^-, Z^0

the force carriers of the fundamental interactions, the vector (*spin 1*) bosons, must be added

Gravitational Graviton

Higgs Boson H^0

scalar (*spin 0*) Higgs boson. The Higgs boson has been observed; it is thought to be the main ingredient in the mechanism that attributes mass to the particles.

The fact that there are so many particles and that so few constitute the present **stable matter** is not currently understood. It is also unclear why the ultimate fermionic constituents appear in **three families**, each constituted of two leptons and two quarks, and each being a replica of the same type, see Table 1



Quantum Numbers of Quarks and Leptons

Table 2 : Leptons and quarks (spin $\frac{1}{2}$ fermions) of the first family

For each particle we have an *anti-particle*: it has the opposite quantum numbers of the *particle* and opposite charge but same mass and spin

		First family			
		Symbol	Q	L_e	B
Leptons	ν_e		0	1	–
	e^-		–1	1	–
Quarks	u		+2/3	–	+1/3
	d		–1/3	–	+1/3

Q is the electric charge in unit of the proton charge, L_e is the electronic lepton number, B the baryonic number.

Stable particles are:

- photon
- neutrinos and the antineutrinos
- the electron, the positron
- proton and the antiproton p

All others are unstable.

- The first family includes the quarks u, d and the leptons e, ν_e .
- The ordinary matter is constituted of quarks u, d and of electrons e .
- The second and third families seem to be “replicas” of the first one. Leptons and quarks of generations higher than first can be produced at accelerators



The Standard Model (2)

Force Carrier	Photon	W & Z Boson	Gluons	Graviton
	EM	Weak	Strong	Gravitational
Quarks	✓	✓	✓	✓
Leptons	✓	✓		✓
Neutrinos		✓		

the neutrinos, being uncharged, are only subject to the weak interaction

The *hadrons* are composed of quarks and are known in two topologies:

- those constituted by three quarks (the *baryons*, like the proton and neutron) and
- those constituted by a quark-antiquark pair (the *mesons*).
- As for leptons, antiquarks also exist and particles composed of three antiquarks are called *antibaryons*.


As will be discussed later, the number of baryons and leptons is conserved (one anti-particle counts with a “-” sign). This means that, as described by the relationship $E = mc^2$, the energy can be converted in mass in the form of particles; nevertheless,

the total number of baryons and leptons must be conserved.

→ If an electron is produced, it must be created in association with a positron (its antiparticle, with electric charge and leptonic number of opposite sign) as expected from the Dirac theory.



Interactions and Vector Bosons



Interactions are mediated by the exchange of **vector bosons**, i.e. particles with spin 1: *photons*, *gluons*, W^+ , W^- and Z^0 bosons. Gravity is mediated by a spin 2 boson, the *graviton*

Graphic representation of the different type of interactions between two particles are shown in the diagrams to the left.

- leptons and quarks by straight lines,
- photons by wavy lines,
- gluons by spirals, and
- W^\pm and Z^0 by dashed lines.

Each of these three interactions is associated with a charge: electric charge, weak charge and strong charge. The strong charge is also called colour charge or colour for short.

A particle is subject to an interaction if and only if it carries the corresponding charge:

- Leptons and quarks carry weak charge.
- Quarks are electrically charged, so are some of the leptons (e. g., electrons).
- Colour charge is only carried by quarks (not by leptons).



Interactions

<i>Vector Boson</i>	<i>Mass</i>	<i>Charge</i>	<i>Comment</i>
Photon	0	N	The rest mass of the photon is zero. Therefore, the range of the electromagnetic interaction is infinite. Photons, however, have no electrical charge → do not interact with each other
W^\pm	≈ 80 GeV/c ²	Y	Heavy particles can only be produced as virtual, intermediate particles in scattering processes for extremely short times. Therefore, the weak interaction is of very short range
Z^0	≈ 91 GeV/c ²	N	
Gluon	0	Y	The gluons, like the photons, have zero rest mass. Gluons, however, carry colour charge. Hence they can interact with each other. As we will see, this causes the strong interaction to be also very short ranged.



Symmetries and Conservation Laws

Symmetries are associated to conservation laws and conservation laws are associated to symmetries.

Effect on a system Ψ of a 'transformation operator' \hat{U}

$$\Psi \rightarrow \psi' = \hat{U}\psi$$

$$(\hat{H}\psi_j = E_j\psi)$$

If physical predictions do not change under $\hat{U} \rightarrow$

- $\hat{U}\hat{U}^\dagger$ must be unitary (if you go & come back you get to the starting point) \rightarrow no change ;
- The operator \hat{U} commutes with the Hamiltonian: $[\hat{H}, \hat{U}] = \hat{H}\hat{U} - \hat{U}\hat{H} = 0$

A continuous transformation (translation or rotation) \sim sequence of 'many' infinitesimal transformations

$$\hat{U}(\epsilon) = \hat{I} + i\epsilon\hat{G}$$

'Infinitesimal transformation \rightarrow transformation ϵ , of 'type' \hat{G}

If $[\hat{H}, \hat{U}] \rightarrow [\hat{H}, \hat{G}] \rightarrow$ An operator acting on a system returns an observable \rightarrow The observable is conserved

(*) or rather the invariance of the equations that describe how a system evolves under transformations



Invariance & Symmetry

physical systems are described by an equation.
The system is considered as invariant if the equation describing it is invariant under given transformation

Important: Invariance properties → conservation laws.

There are two types of transformations:

- **continuous** (like a translation or rotation, it can be seen as a series of infinitesimal transformations)
- **discrete** (like the transition from one state to another).

Symmetry	Conserved Quantity
Spatial Rotation	Angular Momentum
Temporal Translation	Energy
Spatial Translation	Momentum
EM gauge invariance	Electric charge

Classical mechanics: a state with n degrees of freedom is characterised by n q_i coordinates and n conjugated momenta p_i . The evolution of the system is described, in the Lagrangian formalism, by

$$\mathcal{L} = T - V = \text{kinetic energy} - \text{potential energy} \quad p_i = \frac{\partial \mathcal{L}}{\partial \dot{q}_i}$$

$$\frac{dp_i}{dt} - \frac{\partial \mathcal{L}}{\partial q_i} = 0$$

Noether's theorem:
A conserved quantity is associated to a continuous symmetry

If \mathcal{L} does not depend on q_i then $\frac{dp_i}{dt} = 0 \rightarrow$ the conjugated momentum p_i is constant



Continuous Transformations

$$\frac{dp_i}{dt} - \frac{\partial \mathcal{L}}{\partial q_i} = 0$$

• Translation along x

- Let us consider the system Lagrangian $\mathcal{L} = T - V = \frac{1}{2} m \dot{x}^2$. In this case, \mathcal{L} does not depend on x and is invariant under translations along the x axis $\rightarrow p_x = \frac{\partial \mathcal{L}}{\partial \dot{x}} = m\dot{x} = \text{constant}$ i.e., the momentum $p_x = m\dot{x}$ is conserved.
- In Lecture Notes the translation $x \rightarrow x + \epsilon$ inducing $\Psi(x) \rightarrow \psi'(x) = \psi(x + \epsilon)$ in the Hamiltonian representation is also examined to show the same result

• Rotations

- The Lagrangian $\mathcal{L} = T - V = \frac{1}{2} m \dot{\phi}^2 r^2$ where $\dot{\phi} r = v$ does not depend on ϕ ; this implies that \mathcal{L} is invariant under spatial rotations $p_\phi = \frac{\partial \mathcal{L}}{\partial \dot{\phi}} = m\dot{\phi} r^2 = mvr$ is constant, i.e., that the angular momentum is conserved.



Discrete Transformations

Three discrete symmetries are very important in particle physics:

- **spatial parity (P)**: $\psi(\mathbf{r}, t) \rightarrow \psi(-\mathbf{r}, t)$
- **charge conjugation (C)**: $p \rightarrow \bar{p}$
- **time reversal (T)**: $\psi(\mathbf{r}, t) \rightarrow \psi(\mathbf{r}, -t)$

Constraints on possible reactions



Also important

- CP
- CPT

↓

Symmetry Properties based on Observations

Symmetry	Spatial Parity (P)	(C) Charge Conjugation	(T) Time Reversal	(CP)	(CPT)
Strong Interactions	Conserved	Conserved	Conserved	Conserved	Conserved
Elec.magn. Interactions	Conserved	Conserved	Conserved	Conserved	Conserved
Weak Interactions	Not conserved	Not conserved	Conserved	Not always violated	Conserved

All are good symmetries of both the electromagnetic and strong interactions.

- The weak interaction breaks both C and P symmetries maximally but is CP-invariant for many processes. Violation of CP invariance has been observed in the interactions of neutral meson systems, particularly kaons and beauty mesons.
- The product of all three, CPT, is **expected** to be a universal symmetry of physics and is an important assumption of quantum field theory.



Parity of Fermions & Bosons

Bosons and antibosons have integer spin (follow the Bose-Einstein statistics) & the same intrinsic parity. Fermions and antifermions have half integer spin and opposite parities.

It is assumed that

- *Bosons*: $\psi(1,2) = \psi(2,1)$ *symmetric*
- *Fermions*: $\psi(1,2) = -\psi(2,1)$ *anti-symmetric*

$$\begin{aligned}
 &P(\text{quarks/leptons}) = \\
 +1 &= P_{e^-} = P_{\mu^-} = P_{\tau^-} = P_u = P_d = P_s = \dots; \\
 -1 &= P_{e^+} = P_{\mu^+} = P_{\tau^+} = P_{\bar{u}} = P_{\bar{d}} = P_{\bar{s}} = \dots
 \end{aligned}$$

The total wave function of two particles is the product of two terms

Spatial function x **Spin function**

Motion of 1 particle with respect to the other ~ described by spherical harmonics $Y_l^m(\vartheta, \varphi)$

Dirac theory → Spin function is symmetric if the two spins are parallel ↑↑ and anti-symmetric if they are opposite ↑↓

Bosons : both Spatial and Spin functions are symmetric or anti-symmetric

Fermions : one function is symmetric while the other one is anti-symmetric

- $\Psi(x) = \cos(x)$ has positive parity [$\cos(x) = \cos(-x)$]
- $\Psi(x) = \sin(x)$ has negative parity [$\sin(x) = -\sin(-x)$]
- $\Psi(x) = \cos(x) + \sin(x)$ parity is not defined



Symmetries in Particle Physics: Parity

Parity (P) reflection symmetry: depending on whether the sign of the wave function changes under reflection or not, the system is said to have negative or positive **P** respectively. For those laws of nature with left-right symmetry, the parity quantum number **P** of the system is conserved.

Parity refers to a transformation that inverts spatial coordinates, like a reflection:

$$x, y, z \rightarrow -x, -y, -z \quad r \rightarrow -r \quad \text{Exchange right} \leftrightarrow \text{left} \quad [\Theta \rightarrow \pi - \Theta \text{ and } \phi \rightarrow \phi + \pi]$$


$$P\psi(\mathbf{r}) = \psi(-\mathbf{r})$$

If applied twice gives back the original wave function

$$P^2\psi(\mathbf{r}) = PP\psi(\mathbf{r}) = P\psi(-\mathbf{r}) = \psi(\mathbf{r})$$

This implies that eigenvalues of **P** are ± 1 .

- $\Psi(x) = \cos(x)$ has positive parity [$\cos(x) = \cos(-x)$]
- $\Psi(x) = \sin(x)$ has negative parity [$\sin(x) = -\sin(-x)$]
- $\Psi(x) = \cos(x) + \sin(x)$ parity is not defined





Bound States in Atomic Physics: Parity (.. But Why?)

Spatial function x Spin function

An example from atomic physics bound states: *Spin neglected*

$$P_n(x) = (2^n n!)^{-1} \frac{d^n}{dx^n} [(x^2 - 1)^n]$$

$$\psi(r, \theta, \varphi) = \chi(r) Y_\ell^m(\theta, \varphi)$$

ℓ = orbital quantum number

m = azimuthal quantum number

$$Y_\ell^m(\theta, \varphi) = \sqrt{\frac{(2\ell + 1)(\ell - m)!}{4\pi(\ell + m)!}} P_\ell^m(\cos \theta) e^{im\varphi}$$

Spherical harmonics

Legendre polynomials

The space inversion in this case gives: $r \rightarrow -r$, $\Theta \rightarrow \pi - \Theta$ and $\phi \rightarrow \phi + \pi$

Reminder: $\cos(\theta) = -\cos(\pi - \theta)$

If we apply the parity operator P to both Legendre polynomials and to the $e^{im\varphi}$ term P = parity operator

$$P_0(x) = 1$$

$$P_1(x) = x$$

$$P_2(x) = \frac{1}{2}(3x^2 - 1)$$

$$P_3(x) = \frac{1}{2}(5x^3 - 3x)$$

$$P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3)$$

$$P_5(x) = \frac{1}{8}(63x^5 - 70x^3 + 15x)$$

$$P_6(x) = \frac{1}{16}(231x^6 - 315x^4 + 105x^2 - 5)$$

Reminder! Please note that $e^{ix} = \cos(x) + i \sin(x)$


$$P e^{im\varphi} = e^{im(\varphi + \pi)} = e^{im\pi} e^{im\varphi} = (-1)^m e^{im\varphi}$$

$$P P_\ell^m(\cos \theta) = (-1)^{\ell + m} P_\ell^m(\cos \theta)$$

$$P Y_\ell^m(\theta, \varphi) = (-1)^{\ell + m} Y_\ell^m(\theta, \varphi). \quad (-1)^{m+m} = 1$$




$Exp(i\phi)$





Hydrogen Atom (*.. But Why?*)



In 'electric dipole transitions' in the hydrogen atom, $\Delta l = 1$, a photon is emitted

- The parity of the atom changes
- To conserve the parity of the total system (hydrogen atom + photon)

Naming: $J^P = \text{Spin}^{\text{Parity}}$
 Photon $\rightarrow J^P = 1^{-1}$


Photon must have parity -1



Parity Violation in Weak Interactions

The electron neutrino has spin = 1/2 → has (in principle!) two polarization states $\pm 1/2$

Experimental finding: the spin of the electron neutrino is always anti-parallel to the direction of the momentum “left-handed” while the spin of the electron anti-neutrino always points in the same direction of the momentum “right-handed”



Parity transformation changes the momentum but not the spin. If applied to an electron neutrino would create a right-handed electron neutrino

$$P(\vec{p}, \vec{s}) = (\vec{p}, \vec{s})$$

inverts the coordinates

$$\mathbf{r} \Rightarrow -\mathbf{r}$$

does not change time

$$t \Rightarrow t$$

as a consequence

it inverts momenta

$$\mathbf{p} \Rightarrow -\mathbf{p}$$

and does not change angular momenta

$$\mathbf{r} \times \mathbf{p} \Rightarrow \mathbf{r} \times \mathbf{p}$$

including spins

$$\mathbf{s} \Rightarrow \mathbf{s}$$

Since the neutrino only has weak interactions and since the electron neutrino is ONLY left-handed then

Parity is not conserved in weak interactions





Charge Conjugation in Particle Physics

C, charge conjugation, changes particles into antiparticles and vice versa, leaving space coordinates, time and spin unchanged. Therefore, the signs of all the additive quantum numbers, electric charge, baryon number and lepton number are changed.

- Its eigenvalues are ± 1 ; they are multiplicatively conserved in strong and e.m. interactions.
- Only particles (like π^0 , unlike K's) which are their own antiparticles, are eigenstates of \mathbf{C} , with values $C = (\pm 1)$:
C = -1 for γ (accelerated charged particles emit photons; C changes the sign of the particle \rightarrow the photon must compensate!

C = +1 for π^0 ($\pi^0 \rightarrow \gamma\gamma$), η , η' ;

C = -1 for ρ^0 , ω , ϕ ;

- Why define \mathbf{C} ? \rightarrow

C-conservation constraints e.-m. decays:

$\pi^0 \rightarrow \gamma\gamma$: +1 \rightarrow (-1) (-1) ok;

$\pi^0 \rightarrow \gamma\gamma\gamma$: +1 \rightarrow (-1) (-1) (-1) no.

Br($\pi^0 \rightarrow \gamma\gamma\gamma$) measured to be $\sim 10^{-8}$.



Time Reversal


The time reversal operator T reverses the time coordinate t :

$$T t = -t$$

In *classical mechanics*, the systems are invariant under time reversal:

$$T \psi(\mathbf{r}, t) = \psi(\mathbf{r}, -t)$$

- planet moving on a circular orbit around the sun = planet following the same orbit in opposite direction (depending on the initial conditions).
- Similarly, the application of T to a two body scattering process would reverse the reaction:



If the parity P (or T) is conserved \rightarrow the Hamiltonian must not contain terms that change sign due to P or T .

\rightarrow particles with spin may have a magnetic dipole moment, but not a static electric dipole moment (as in ordinary matter) because the term is not invariant under T .

Quantity	Transformation		
	P	T	
\mathbf{r}	$-\mathbf{r}$	\mathbf{r}	Polar vector
\mathbf{p}	$-\mathbf{p}$	$-\mathbf{p}$	Polar vector
σ	σ	$-\sigma$	Axial vector (like $\mathbf{L} = \mathbf{r} \times \mathbf{p}$)
\mathbf{E}	$-\mathbf{E}$	\mathbf{E}	Remember that $\mathbf{E} = -\partial\phi/\partial\mathbf{r}$
\mathbf{B}^a	\mathbf{B}	$-\mathbf{B}$	\mathbf{B} is similar to σ

^aWe can think of \mathbf{B} as being due to the current in a coil. Reversing T means to invert the current direction and therefore the magnetic field direction



Invariance Properties of Fundamental Interactions

Additive (A) or Multiplicative (M) quantum number

Continuous Symmetries



Discrete Symmetries


Conservation of	Interaction			<i>N</i>
	Strong	Electromagnetic	Weak	
Energy–Momentum $E; \mathbf{p}$	yes	yes	yes	A
Angular momentum \mathbf{J}	yes	yes	yes	A
Parity P	yes	yes	no	M
Baryonic number B	yes	yes	yes	A
Leptonic numbers ^b L_e, L_μ, L_τ	yes	yes	yes	A
Electric charge Q	yes	yes	yes	A
Charge conjugation C	yes	yes	no	M
Time reversal T	yes	yes	yes ^a	M
CP	yes	yes	yes ^a	M
CPT	yes	yes	yes	M
Strong isospin I	yes	no	no	A
3 rd isospin component I_z	yes	yes	no	A
Strangeness S	yes	yes	no	A
Lifetime	$\sim 10^{-23}$ s	$\sim 10^{-20}$ s	$\sim 10^{-12}$ s	–
Interaction range	$\sim 10^{-13}$ cm	infinite	$< 10^{-15}$ cm	–

Will see later!



CP and CPT

Weak interactions violate C and P. However they were (believed to be) invariant under the CP transformation



In '60 a rare decay of a neutral Kaon was discovered which was violating CP. A very small effect (\rightarrow rare decay!) \rightarrow important consequence in the evolution of the Universe:

- At the time of Big Bang the number of particles = the number of anti-particles
- Due to CP violation the number of particles became slightly larger than the number of anti-particles
- Annihilations left only particles in excess \rightarrow Universe is made of particles

However the invariance under CPT transformations is a fundamental property of quantum field theories
 \rightarrow the CP violation also involves a violation of T to maintain this symmetry



Isospin, a new Quantum Number

- $m_p \approx m_n ; \sigma_{pp} \approx \sigma_{pn}$
- If you exchange $n \leftrightarrow p$ in Nuclei they remain very similar: ${}^7\text{Li}(3p + 4n) \approx {}^7\text{Be}(4p + 3n)$, ${}^{13}\text{C}(6p + 7n) \approx {}^{13}\text{N}(7p + 6n)$

In ~1930 Heisenberg, Condon and Carren made the hypothesis:
 proton and the neutron are *two different states of the nucleon*.

→ A new quantum number: the *Isospin*, I . The nucleon was chosen to have $I=1/2 \rightarrow I_3$ projections:
 the proton (+1/2) and the neutron (-1/2)

baryons	$m(\text{MeV}/c^2)$	B	Q	S	mesons	$m(\text{MeV}/c^2)$	B	Q	S
p	938.272	+1	+1	0	K^+	493.68	0	+1	+1
n	939.565	+1	0	0	K^0	497.65	0	0	+1
Λ	1115.68	+1	0	-1	η	547.7	0	0	0
Σ^+	1189.4	+1	+1	-1	π^+	139.570	0	+1	0
Σ^0	1192.6	+1	0	-1	π^0	134.977	0	0	0
Σ^-	1197.4	+1	-1	-1	π^-	139.570	0	-1	0
Ξ^0	1314.8	+1	0	-2	\bar{K}^0	497.65	0	0	-1
Ξ^-	1321.3	+1	-1	-2	K^-	493.68	0	-1	-1

Isospin combines like the spin:
 for a value I of the Isospin you
 have $(2I + 1)$ possible
 combinations $\rightarrow I_3$ values.
 $0 \rightarrow$ singlet
 $1/2 \rightarrow$ doublet
 $1 \rightarrow$ triplet
 The state np ($I_3=+1/2-1/2$) may
 belong to a singlet or triplet



Isospin and Symmetry of Wave Functions

A rotation in the space of the Isospin does not change the state, Isospin is a conserved quantity in strong interactions (but not in weak and electro-weak). Mass differences are only due to EM interactions

If you have a nucleus with Z protons and N neutrons the projection of the Isospin will be

$$I_3 = \frac{1}{2}Z - \frac{1}{2}N = \frac{(Z-N)}{2}$$

Considering that the baryon number is $B = Z + N$ and that the charge is $Q = Z$ one can write

$$Q = I_3 + B/2$$

Isospin has to be considered when studying the symmetry of a pair of fermions:

$$\Psi = \psi(\text{Space})\chi(\text{spin})I(\text{Isospin})$$

$$\text{Space} \rightarrow -1^L$$

$$\text{Spin} \rightarrow -1^{S+1}$$

$$\text{Isospin} \rightarrow -1^{I+1}$$

The symmetry of a system with L, S, I goes like

$$\text{Symmetry} \rightarrow (-1)^L (-1)^{S+1} (-1)^{I+1}$$

A system with two nucleons has to be anti-symmetric as requested by the Pauli principle



Mesons Isospin

baryons	$m(\text{MeV}/c^2)$	B	Q	S	mesons	$m(\text{MeV}/c^2)$	B	Q	S
p	938.272	+1	+1	0	K^+	493.68	0	+1	+1
n	939.565	+1	0	0	K^0	497.65	0	0	+1
Λ	1115.68	+1	0	-1	η	547.7	0	0	0
Σ^+	1189.4	+1	+1	-1	π^+	139.570	0	+1	0
Σ^0	1192.6	+1	0	-1	π^0	134.977	0	0	0
Σ^-	1197.4	+1	-1	-1	π^-	139.570	0	-1	0
Ξ^0	1314.8	+1	0	-2	\bar{K}^0	497.65	0	0	-1
Ξ^-	1321.3	+1	-1	-2	K^-	493.68	0	-1	-1

$$Q = I_3 + B/2$$

Protons and neutrons we saw already, all OK

Pion's masses are close by and may be considered as members of the same triplet with $l=1$ and $l_3=-1,0,1$. Also the charges are correctly computed using the standard formula (baryon number=0)

The η has a mass very different from the pion's mass and it is ~isolated \rightarrow only member of a singlet. Charge is OK, $l=0, l_3=0$

K^+K^0 are also close in mass, like the pair $K^-\bar{K}^0$, may be assumed to be members of a doublet, $l=1/2, l_3=-1/2, 1/2$. However the Q formula fails.

All is restored if we include S, the *strangeness*, in the charge formula and define a new quantum number, the

$$\text{Hypercharge } Y = B + S$$

$$Q = I_3 + \frac{B + S}{2} = I_3 + \frac{Y}{2}$$



Baryon Isospin

baryons	$m(\text{MeV}/c^2)$	B	Q	S	mesons	$m(\text{MeV}/c^2)$	B	Q	S
p	938.272	+1	+1	0	K^+	493.68	0	+1	+1
n	939.565	+1	0	0	K^0	497.65	0	0	+1
Λ	1115.68	+1	0	-1	η	547.7	0	0	0
Σ^+	1189.4	+1	+1	-1	π^+	139.570	0	+1	0
Σ^0	1192.6	+1	0	-1	π^0	134.977	0	0	0
Σ^-	1197.4	+1	-1	-1	π^-	139.570	0	-1	0
Ξ^0	1314.8	+1	0	-2	\bar{K}^0	497.65	0	0	-1
Ξ^-	1321.3	+1	-1	-2	K^-	493.68	0	-1	-1


$$Q = I_3 + \frac{B + S}{2} = I_3 + \frac{Y}{2}$$

Protons and neutrons we saw already, all OK

Baryons seem to be organised into multiplets as mesons: 1 singlet, 2 doublets, 1 triplet.
The charge-formula works well for baryons!



Extension to Other Generations ($\rightarrow c, b, t$ quarks)



Generalised charge formula $Q = I_z + \frac{B + S + C + B + T}{2}$

Baryon number \downarrow Y

Bottomness number \uparrow

Property / Quark	d	u	s	c	b	t
Q – electric charge	$-\frac{1}{3}$	$+\frac{2}{3}$	$-\frac{1}{3}$	$+\frac{2}{3}$	$-\frac{1}{3}$	$+\frac{2}{3}$
I – isospin	$\frac{1}{2}$	$\frac{1}{2}$	0	0	0	0
I_z – isospin z-component	$-\frac{1}{2}$	$+\frac{1}{2}$	0	0	0	0
S – strangeness	0	0	-1	0	0	0
C – charm	0	0	0	+1	0	0
B – bottomness	0	0	0	0	-1	0
T – topness	0	0	0	0	0	+1



Life-time of Particles (.. the path to Resonances)

Stability of particles:

Stable particles: (...believed to be...) are:

- the photon γ ,
- the electron e^- ,
- neutrinos ν .
- the proton (and the antiproton) (only stable hadron!)

} (and the corresponding antiparticles)

In some models 'Beyond Standard Model' BSM also the proton and the neutrinos may unstable.

Interaction	Life-time (s)	Comment	Example
Weak	10^{-6} to 10^{-12}	~ directly measurable	$\mu^- \rightarrow e^- \bar{\nu}_e \nu_\mu$
Electro-magnetic	10^{-16} to 10^{-20}		$\pi^0 \rightarrow \gamma\gamma$
Hadronic	10^{-23}	'Resonances' (more in the following!)	$\Lambda \rightarrow p\pi^-$
W^+, W^-, Z^0	$\sim 10^{-25}$	Decay is very fast due to large mass \rightarrow light particles	$Z^0 \rightarrow \mu^+ \mu^-$

How to measure these short filetimes?



Of the Uncertainty Principle

Sylvie Braibant, Giorgio Giacomelli, Maurizio Spurio:
Particles and Fundamental Interactions

Clarifications regarding the uncertainty principle. The uncertainty principle tells us that in Nature, a limit exists on our possible knowledge of the submicroscopic world, e.g., regarding the dynamics of a particle. For pairs of conjugated physics variables, for example, energy and time, momentum and position, there are limitations in the precision of their measurements. For example, if we measure the position x of an electron with a precision Δx , we cannot simultaneously measure the p_x component of its momentum with unlimited precision. According to the uncertainty principle, an uncertainty Δp_x related to the uncertainty Δx exists. Similarly, ΔE and Δt are related through the uncertainty principle. In the literature, different numerical expressions for the uncertainty principle are used, that is,

$$\Delta E \Delta t \geq \frac{\hbar}{2}, \quad \Delta E \Delta t \geq \hbar, \quad \Delta E \Delta t \geq h$$

$$\Delta p_x \Delta x \geq \frac{\hbar}{2}, \quad \Delta p_x \Delta x \geq \hbar, \quad \Delta p_x \Delta x \geq h.$$

$\Delta t, \Delta E, \Delta x, \Delta p$: a measurement repeated many times will be in 67% of cases 'within' $1 \Delta t, \Delta E, \Delta x, \Delta p$

Resonance (more in the following!):
 Δt , Lifetime τ
 ΔE , Width Γ
 $\rightarrow \tau \Gamma \geq \hbar$

Undergraduate Lecture Notes in Physics

Sylvie Braibant
Giorgio Giacomelli
Maurizio Spurio

Particles and Fundamental Interactions

An Introduction to Particle Physics

Second Edition

Springer




Resonances and the Uncertainty Principle

The *Uncertainty Principle* relates pairs of conjugated physics variables:
(energy and time), (position and momentum)

$$\Delta E \Delta t \geq \frac{\hbar}{2}$$

$$\Delta p_x \Delta x \geq \frac{\hbar}{2}$$

interaction of two particles




R: unstable \rightarrow decays

Invariant mass
 $a' + b' = m_R$

R: possible intermediate state

Peak at the resonance
mass



In the case of a resonance, we consider the energy (mass) and width Γ (uncertainty on the energy \rightarrow on the mass), lifetime of the particle at rest τ .

$$\Delta E \sim \Gamma \quad \text{The width can be measured!}$$

$$\Delta t \sim \tau \quad \text{Too short to be measured!}$$

$$\rightarrow \Delta E \Delta t = \Gamma \tau \geq \frac{\hbar}{2}$$

- For a $\Gamma \sim 100 \text{ MeV} \rightarrow \tau \sim 10^{-23} \text{ s}$
- If a resonance is produced with an energy $E \rightarrow$ lifetime is increased by a factor $\gamma = E/m \rightarrow \tau \gamma$




Hadronic Resonances

- a particle with $\tau \geq 10^{-12} \text{ s}$ (\rightarrow weak interactions) travels enough to be detected.
- Hadron resonances \rightarrow strong interactions \rightarrow their life-time is so short that it cannot be measured.

How to get their life-time? $\rightarrow \Delta E \Delta t = \Gamma \tau \geq \frac{\hbar}{2} \rightarrow$ repeated measurements of the mass will give different results.

The width of the distribution is connected to the life-time: $\Gamma = \Delta M c^2 = \frac{\hbar}{2} / \tau$

Cross section maximal at the peak of the resonance



Measured resonance mass (inv.mass final state particles)

(Will derive this expression soon)

$$|\chi(Mc^2)|^2 \propto \frac{1}{(Mc^2 - M_0c^2)^2 + \frac{\Gamma^2}{4}}$$

Measured mass value Resonance mass value Resonance width Γ

A width of ~ 100 MeV corresponds to a life-time of 10^{-23} s \rightarrow much too short to be measured!

The presence of a resonance is signalled by increase of cross section !!!!




Getting the Breit Wigner Shape

Intermediate excited states may show up in hadronic interactions. These states R are called resonances. \rightarrow described by a wave function with a de Broglie $\lambda = \frac{h}{p} = \frac{2\pi\hbar}{pc}$ and described by a wave-function like $\psi(t) = \psi(0)e^{-\frac{iE_R}{\hbar}t}$. This state is unstable and will decay to a' and b' \rightarrow **decay law**

Elastic scattering case \rightarrow same particles of the initial state also in the final state (but different momenta)

Ideal Experiment:


Beam a of variable energy


Target containing b (at rest)



If you increase the energy of particle a an intermediate state R may be produced \rightarrow increase in the cross section.

dP = probability of decay per unit time


N = number of produced resonances

$\rightarrow -dN = \lambda N dt$ where λ is a constant that describes how quickly the resonance decays.

$$N(t) = N_0 e^{-\lambda t} = N_0 e^{-t/\tau}$$



Getting the Breit Wigner Shape




Scattering \rightarrow 2 particles in the final state

Elastic scattering \rightarrow same particles of the initial state

1 $\psi(0)e^{-i\omega_R t}$
free particle wave function

2 $N(t) = N_0 e^{-t/\tau}$
decay probability vs time

$$\psi(t) = \psi(0)e^{-i\omega_R t} \cdot e^{-t/2\tau} = \psi(0)e^{-iE_R t/\hbar} \cdot e^{-t \cdot \Gamma/2\hbar} \quad (\text{where } \omega_R = \frac{E_R}{\hbar} \text{ and } \tau = \frac{\hbar}{\Gamma})$$



The probability $I(t)$ of finding the resonance decay at a time t is

$$I(t) = \psi^* \psi = \psi(0)^2 \cdot e^{-\frac{t}{\tau}}$$

$$I(t) = I(0) \cdot e^{-\frac{t}{\tau}} \quad \text{Exponential life-time}$$

Which is the probability $\chi(E)$ of producing a resonance with energy E ? \rightarrow Do a Fourier transform

The Fourier transform is a transformation that decomposes functions depending on space or time into functions depending on spatial or temporal frequency \rightarrow in our case (resonance) gives us the energy distribution

$$\chi(E) = \int \psi(t) e^{iEt} dt = \psi(0) \int e^{-iE_R t/\hbar} \cdot e^{-\Gamma/2\hbar} e^{iEt} dt$$



Relation between Width and Cross Section

$$\chi(E) = \int \psi(t) e^{iEt} dt = \psi(0) \int e^{-iE_R t/\hbar} \cdot e^{-\Gamma/2\hbar} e^{iEt} dt$$

(Let's remember that $\int_0^\infty e^{-ax} dx = \frac{1}{a}$)

$$\rightarrow \chi(E) = \frac{\text{Constant}}{(E_R - E) - i\Gamma/2}$$

We have put $\hbar = 1$

Maximum at $E=E_R$

$$\sigma(E) = \sigma_0 \cdot \chi^*(E) \cdot \chi(E) = \sigma_0 \cdot \frac{\text{Constant}^2}{[(E_R - E)^2 + \Gamma^2/4]}$$

The *Constant* has to be related to the cross section: the square of the wave function $\chi(E)$ represents the probability of finding the particle in the energy state E, it must be proportional to the process cross-section, that is

$$\sigma_0 \sim \text{wave - length}^2 \sim 4\pi\lambda^2 \propto 1/p^2$$


$$\sigma(E) \sim 4\pi\lambda^2 \cdot \frac{\text{Constant}^2}{[(E_R - E)^2 + \Gamma^2/4]}$$

Maximum at $E_R \rightarrow$ we can write

$$1 = \chi^*(E_R) * \chi(E_R) = \frac{\text{Constant}^2}{\Gamma^2/4} \rightarrow \text{Constant}^2 = \frac{\Gamma^2}{4}$$

Breit – Wigner formula (sometimes used as synonym of ‘resonance’)


$$\sigma(E) = 4\pi\lambda^2 \frac{\Gamma^2/4}{(E_R - E)^2 + \Gamma^2/4}$$





Taking Spin into account

Resonance with spin J produced by the collision of two particles a, b , with spin s_a, s_b



Spin sub-states of the resonance: $(2J + 1)$

Spin sub-states the initial state: $(2s_a + 1)(2s_b + 1)$

→ The cross-section must be averaged over the number of spin states of the incoming particles and multiplied by a factor $(2J + 1)$

$$\sigma(E) = 4\pi\lambda^2 \frac{(2J + 1)}{(2s_a + 1)(2s_b + 1)} \frac{\Gamma^2/4}{(E_R - E)^2 + \Gamma^2/4}$$

- The measurement of the cross section also allows to constraint the spin of the colliding particles.
- Spins of the incoming particles are known → Factor $(2J+1)$ gives the spin of the resonance




One observation for inelastic collisions

What we said is valid in the elastic case when particles in the final state are the same of the initial state




$$\sigma(E) = 4\pi\lambda^2 \frac{(2J + 1)}{(2s_a + 1)(2s_b + 1)} \frac{\Gamma^2/4}{(E_R - E)^2 + \Gamma^2/4}$$



Elastic case

Γ^2 is ok for the elastic case because you have the same particles in the initial and final state
 Γ tells us how strongly R couples to (ab) once in the initial state and once in the final state

$$\rightarrow \Gamma_{ab} \cdot \Gamma_{ab} \rightarrow \Gamma^2$$



X Inelastic case

In the inelastic case $a + b \rightarrow R \rightarrow X$ ('many particles')

$$\rightarrow \Gamma_{ab} \cdot \Gamma_X$$


In the inelastic case you may have different decays \rightarrow define different 'Branching fractions'



Resonance : σ_R Inelastic Case

$$\sigma_0 \sim \text{wave-length}^2 \quad \lambda = \frac{h}{p} = \frac{2\pi\hbar}{pc}$$

$$\rightarrow s \propto p_{a,b}$$



(E, \vec{p}) : CM 4-mom.
 Γ_R : constant width
 $\Gamma_{ab, X}$: couplings
 M_R : E_0 , mass

$$\sigma_{ab \rightarrow R \rightarrow X}(E_{\text{CM}} = \sqrt{s}) = \frac{\pi}{|\vec{p}_{a,b}|^2} \frac{(2J_R + 1)}{(2S_a + 1)(2S_b + 1)} \frac{\Gamma_{ab}\Gamma_X}{(\sqrt{s} - M_R)^2 + \Gamma_R^2/4} \approx$$


$$\approx \left[\frac{16\pi}{s} \right] \left[\frac{(2J_R + 1)}{(2S_a + 1)(2S_b + 1)} \right] \left[\frac{\Gamma_{ab}}{\Gamma_R} \right] \left[\frac{\Gamma_X}{\Gamma_R} \right] \left[\frac{\Gamma_R^2/4}{(\sqrt{s} - M_R)^2 + \Gamma_R^2/4} \right]$$

scale factor
(1/s)


statistical factor
(particle spins)

= BR(R \rightarrow ab)

= BR(R \rightarrow X)



$$\sigma(e^+e^- \rightarrow J/\psi \rightarrow \mu^+\mu^-) = \left[\frac{16\pi}{s} \right] \left[\frac{3}{4} \right] \left[\frac{\Gamma_{ee}}{\Gamma_{\text{tot}}} \right] \left[\frac{\Gamma_{\mu\mu}}{\Gamma_{\text{tot}}} \right] \left[\frac{(\Gamma_{\text{tot}}/2)^2}{(\sqrt{s} - M)^2 + (\Gamma_{\text{tot}}/2)^2} \right] =$$



$$= \frac{12\pi}{s} \text{BR}_{J/\psi \rightarrow e^+e^-} \text{BR}_{J/\psi \rightarrow \mu^+\mu^-} \left[\frac{(\Gamma_{\text{tot}}/2)^2}{(\sqrt{s} - M)^2 + (\Gamma_{\text{tot}}/2)^2} \right]$$

$e^+e^- \rightarrow J/\psi \rightarrow \mu^+\mu^-$
 $\sigma_{\text{peak}} \propto 1/s$ ($\approx M_R^{-2}$),
independent from coupling strength.



Resonance : Different Functions

Many more parameterizations used in literature (semi-empirical or theory inspired), e.g.:

$$\sigma_0 = \left[\frac{16\pi}{(2p)^2} \right] \left[\frac{(2J_R + 1)}{(2S_a + 1)(2S_b + 1)} \right] \left[\frac{\Gamma_{ab}}{\Gamma_R} \right] \left[\frac{\Gamma_{final}}{\Gamma_R} \right] \left[\frac{\Gamma_R^2/4}{(\sqrt{s} - M_R)^2 + \Gamma_R^2/4} \right]$$

original, non-relativistic

$$\sigma_1 = \left[\frac{16\pi}{s} \right] \left[\frac{(2J_R + 1)}{(2S_a + 1)(2S_b + 1)} \right] \left[\frac{\Gamma_{ab}}{\Gamma_R} \right] \left[\frac{\Gamma_{final}}{\Gamma_R} \right] \left[\frac{\Gamma_R^2/4}{(\sqrt{s} - M_R)^2 + \Gamma_R^2/4} \right]$$

$m_a, m_b \ll p$

$$\sigma_2 = \left[\frac{16\pi}{M_R^2} \right] \left[\frac{(2J_R + 1)}{(2S_a + 1)(2S_b + 1)} \right] \left[\frac{\Gamma_{ab}}{\Gamma_R} \right] \left[\frac{\Gamma_{final}}{\Gamma_R} \right] \left[\frac{\Gamma_R^2/4}{(\sqrt{s} - M_R)^2 + \Gamma_R^2/4} \right]$$

simpler, neglect s-dependence

$$\sigma_3 = \left[\frac{16\pi}{M_Z^2} \right] \left[\frac{3}{4} \right] \left[\frac{\Gamma_{ee}}{\Gamma_Z} \right] \left[\frac{\Gamma_{ff}}{\Gamma_Z} \right] \left[\frac{M_Z^2 \Gamma_Z^2}{(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} \right]$$

relativistic BW for $e^+e^- \rightarrow Z \rightarrow ff$





$$\sigma_4 = \left[\frac{16\pi}{M_Z^2} \right] \left[\frac{3}{4} \right] \left[\frac{\Gamma_{ee}}{\Gamma_Z} \right] \left[\frac{\Gamma_{ff}}{\Gamma_Z} \right] \left[\frac{s \Gamma_Z^2}{(s - M_Z^2)^2 + s^2 \Gamma_Z^2 / M_Z^2} \right]$$

"s-dependent Γ_Z " (used at LEP for the Z lineshape)

Resonances: Practicalities

There are two mechanisms for the observation of resonances:


- **Formation:** the two interacting particles have both quantum numbers and energy to produce a resonance. \rightarrow increase of the cross-section at an energy corresponding to $M_R \rightarrow$ mass & width \rightarrow life-time \rightarrow **EASY!**
- **Production:** the two interacting particles do **NOT** have the quantum numbers to create a resonance \rightarrow an intermediate virtual particle is needed (with correct quantum numbers!). Determining the presence of this resonance will be more **DIFFICULT** \rightarrow
 - identify decay products of R (use 2 right particles out of 3)
 - construct of the invariant mass.





The Electromagnetic Paradigm

- **Electromagnetic (EM):** The analytic form of the interaction potential between charged particles is precisely known → Maxwell's original formulation → relativistic representation → quantized field theory.
- **Quantum electrodynamics (QED):** includes the spin of particles, the interaction between charged particles through the exchange of a photon. Many physics quantities (cross-sections, particle lifetimes, magnetic moments, and so on) can be computed very precisely.
- QED extended to the weak and (partially) to the strong interactions → *comparison of theoretical predictions with experimental measurements*.
- electromagnetic and weak interactions are different manifestations of a single interaction, the electroweak interaction. The **unification** of the two interactions occurs at an energy $\sim W^+, W^-$, and Z^0 masses ~ 100 GeV. At lower energies, the electromagnetic and weak interactions are separate and different.
- At much larger energies, the electroweak and strong interaction unification (the so-called **Grand Unification Theory**) can be hypothesized.





The Electro-Magnetic Case

Coulomb force:


$$F = K \frac{q_1 q_2}{r^2} \vec{r}$$

- q_1 e q_2 are the point-like particle electric charges,
- r is the distance between them,
- \vec{r} is a unit vector directed from q_1 to q_2 (positive or negative) and
- K is a proportionality constant. The dependence on r is similar to that of Newton's law.
- The *electrostatic force can attract or repel particles*, depending on the relative sign of the charges.

Magnetic field is generated by electric charges in motion; the force acting on a moving charge q with velocity \mathbf{v} in an electric field \mathbf{E} and a magnetic field \mathbf{B} is:

$$F = q\mathbf{E} + q\mathbf{v} \times \mathbf{B}. \quad \text{Classical approach: interaction between particle \& field}$$


Today's view: Graphic representation (\rightarrow Feynman diagrams) for the elastic electron–electron interaction.



- (a), the electron in the bottom part emits a “virtual” photon which is then absorbed by the electron at the top;
- (b) the other way around.
- (c) the interaction without specifying who emits the photon



The EM Case & Feynman Diagrams



- Feynman diagrams have been very successful for describing the EM interactions.
 - both an intuitive representation of the interaction and
 - a rigorous way to obtain numerical quantities through a perturbative calculation method (see later)

The interaction between two electrons: they repel each other.
The interaction happens through the exchange of a photon (in this case).

An electron at **rest** cannot, however, emit a “real” photon because this would violate the energy conservation law


Process	Initial state energy	Final state energy	
$e \rightarrow e\gamma$	$m_e c^2$	$\neq m_e c^2 + \frac{p_e^2}{2m_e} + E_\gamma$	(non relativistic case)

E_γ is the total energy of the emitted photon, p_e is the (nonrelativistic) momentum acquired by the electron, m_e is the electron mass. According to Heisenberg’s uncertainty principle, if energy is measured with an uncertainty of ΔE , the uncertainty on the time measurement is

$$\Delta t \geq \hbar / (\Delta E).$$



The Nature of the Electromagnetic Interaction



No solution??? Let's follow the evolution of the process:

$$\Delta t \geq \hbar / (\Delta E).$$

- A photon is emitted from the first electron violating the energy conservation (by a quantity ΔE).
- The photon is absorbed by the second electron after a time t , \rightarrow a second violation of energy conservation by a value of $-\Delta E$.
- If all this happens within Δt of the uncertainty principle, *the two violations cannot be observed*: they are “hidden” by the uncertainty principle.

This process is considered as possible.

The net effect is an exchange of energy and momentum between the two electrons, and is therefore a way in which two electrons, and more generally two charged particles, can interact.

The electron quantum numbers, particularly its spin, must remain unchanged. \rightarrow

As a consequence, the exchanged particle must have integer spin and is therefore a boson (all the force mediators have spin 1, except the graviton that has spin 2).



The Nature of the Electromagnetic Interaction

The photon is massless \rightarrow moving at the light speed, it travels in the time interval Δt a distance

$$\Delta r = c\Delta t$$

Placing this quantity in the uncertainty relation, one obtains

$$\Delta E \geq \hbar/(\Delta t) \approx \hbar c/(\Delta r).$$


Since the interaction energy V is of the order of ΔE one has

$$\Delta E \simeq V = \alpha_i \hbar c/r.$$

The dimensionless constant α_i gives the interaction intensity \rightarrow the forces due to the exchange of virtual massless particles decrease with the distance r as

$$F \sim dV/dr \sim 1/r^2$$

- From the opposite approach, $1/r^2 \rightarrow$ the exchanged virtual particle is massless.
- Since the gravitational force has a similar dependence in $1/r^2$, the graviton should also be massless.



Comment:

Since the virtual field (the photon) is not observable, and since the final states are identical, we do not know if the photon is created (destroyed) by the electron at the top or at the bottom of the diagram: the two processes are indistinguishable.



Elaborating more

The dimensionless parameter characteristic of the EM interaction is the fine structure constant (also called electromagnetic coupling constant) already known from atomic physics. It can be derived by equating

$$\Delta E \simeq V = \alpha_i \hbar c / r.$$

with the Coulomb energy potential:

$$\frac{\alpha_i \hbar c}{r} = K q^2 / r$$

from which one finds ($q = e$ is the electric charge of the electron):

$$\alpha_i = e^2 / \hbar c$$

numerically, one has (S.I., cgs, cgs)

electromagnetic coupling constant

$$\mathbf{F}_1 = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r_{12}^2} \mathbf{e}_{12} = -\mathbf{F}_2.$$

$$\alpha_{EM} = \frac{e^2}{4\pi\epsilon_0 \hbar c} = \frac{(1.602 \cdot 10^{-19})^2}{4\pi \cdot 8.85 \cdot 10^{-12} \cdot 1.05 \cdot 10^{-34} \cdot 3 \cdot 10^8} = \mathbf{1/137.1}$$

$$\alpha_{EM} = \frac{e^2}{\hbar c} = \frac{(4.803 \cdot 10^{-10})^2}{1.0546 \cdot 10^{-27} \cdot 3 \cdot 10^{10}} = \mathbf{1/137.1} = 7.294 \cdot 10^{-3}$$


$$\alpha_{EM} = \mathbf{e^2} \quad (\hbar = c = 1)$$

1/137 is a ~small number \rightarrow
the perturbation induced by the
exchange of one photon is ~small

Will discuss more about it



Micro to Macro world




No way to 'see' what is in the microscopic world → can only see the effect of sending a projectile on your target



Nuclear Sizes → why electron scattering?

Nuclear sizes and shapes → use scattering technique → use a projectile (accelerated or from radioactivity) that hits a target

Protons are extended and complex objects



nuclear forces between the projectile and the target are complex and not well understood

Use electrons! Point-like projectiles!

- *The interactions between an electron and a nucleus, nucleon or quark takes place via the exchange of a virtual photon — this may be very accurately calculated in quantum electrodynamics (QED).*
- *These processes are in fact manifestations of the well known electromagnetic interaction, whose coupling constant $\alpha \approx 1/137$ is much less than one. This last means that higher order corrections play only a tiny role*



Preparing electron scattering studies!

In electron scattering experiments one employs highly relativistic particles \rightarrow use four-vectors in calculations. The zero component of space-time four-vectors is *time*, the zero component of four momentum vectors is *energy*:

$$x = (x_0, x_1, x_2, x_3) = (ct, \mathbf{x}),$$
$$p = (p_0, p_1, p_2, p_3) = (E/c, \mathbf{p}).$$

Three-vectors are '**bold**'. The Lorentz-invariant scalar product of two four-vectors a and b is defined by

$$a \cdot b = a_0 b_0 - a_1 b_1 - a_2 b_2 - a_3 b_3 = a_0 b_0 - \mathbf{a} \cdot \mathbf{b}.$$

Let's compute the four-momentum squared $p^2 = \frac{E^2}{c^2} - \mathbf{p}^2$.

This squared product is equal to the square of the rest mass $m^2 c^2$. In fact it is always possible to find a reference frame in which the particle is at rest $\rightarrow \mathbf{p} = \mathbf{0}$, and $E = mc^2$. The quantity

$$m = \sqrt{p^2}/c \text{ is called the invariant mass}$$

From the two relations above we obtain $E^2 - \mathbf{p}^2 c^2 = m^2 c^4$

for electrons \sim high energy electrons (already at energies of a few MeV) $E \approx |\mathbf{p}| c$ if $E \gg mc^2$.



The 'low energy' case

$$\begin{aligned} E^2 &= m^2 + p^2 \quad (c = 1) \\ E &= m \cdot \sqrt{1 + \frac{p^2}{m^2}} \\ &\approx m \cdot \left(1 + \frac{1}{2} \frac{p^2}{m^2} + \dots \right) \approx m + \frac{p^2}{2m} \\ &\approx \frac{p^2}{2m} \end{aligned}$$

To answer, let us consider $v \ll c$.

Then $\vec{p} \approx \vec{v} \frac{E_0}{c^2} = m\vec{v}$

and $E = E_0 + T = \sqrt{p^2 c^2 + m^2 c^4}$

$$\begin{aligned} &= mc^2 \left(1 + \frac{p^2 c^2}{m^2 c^4} \right)^{1/2} \\ &\approx mc^2 \left(1 + \frac{1}{2} \frac{p^2 c^2}{m^2 c^4} + \dots \right) \\ &= mc^2 + \frac{p^2}{2m} + \dots \end{aligned}$$



(Geometric) Cross Sections – 1 (Povh...)


Structure of the matter is studied with scattering experiments. Energetic projectiles → small equivalent wave length

$$\lambda = \hbar/p$$

Ideal Simplified Experiment:

Beam particles **a** bombard scattering centres **b**.


- reaction occurred when **a** hits **b**.
- The beam particle **a** disappears after the interaction



Particle beam **a** coming from left with density n_a and velocity v_a . The corresponding flux is

$$\phi_a = n_a \times v_a$$

Target with N_b scattering centres **b** and particle density n_b



$$N_b = A d n_b$$

$$= (\text{density} \times \text{Volume})_{\text{target}}$$



(Geometric) Cross Sections - 2 (Povh...)

Ideal Simplified Experiment:

After the interaction beam particles disappear (we do not distinguish different final topologies, we sum elastic + inelastic cross sections). Reaction rate is

$$\dot{N} = N_a - \dot{N}_a$$

Particle beam **a** coming from left with density n_a and velocity v_a . The corresponding flux is

$$\phi_a = n_a \times v_a = \frac{N_a}{A} (\text{area} \times \text{time})^{-1}$$

Target with N_b scattering centres **b** and particle density n_b .

Target particles within the beam area A are

$$N_b = A \times d \times n_b$$


→ the reaction rate \dot{N} is

$$\dot{N} = \phi_a \times N_b \times \sigma_b$$

$$\sigma_b = \frac{\dot{N}}{\phi_a \times N_b}$$

number of reactions per unit time

beam particles per unit time per unit area × scattering centres



Limitations: HP, scattering centres do not overlap + only one scattering



(Geometric) Cross Sections - 3 (Povh...)

If beam is not uniform

$$\sigma_b = \frac{\dot{N}}{\phi_a \times N_b} = \frac{\text{number or reactions per unit time}}{\text{beam particles per unit time} \times \text{scattering centres per unit area}}$$

In the expression

$$\sigma_b = \frac{\dot{N} \text{ Physics!}}{\phi_a \times N_b \text{ Experiment}}$$

$(\phi_a \times N_b) =$ Luminosity, \mathcal{L} in this case

- Energy dependence
- Particle types..

$$\dot{N} = \mathcal{L} \times \sigma_b$$

The total cross section σ_{tot} is as the sum of elastic and inelastic cross section

$$\sigma_{tot} = \sigma_{el} + \sigma_{inel}$$

and has dimensions of area. a common unit to define cross sections is the **barn**

$$\sigma_{pp}(10 \text{ GeV}) \sim 40 \text{ mb}, \sigma_{vp}(10 \text{ GeV}) \sim 70 \text{ fb} \text{ (ratio is } \rightarrow 10^{-12})$$

Unit	Symbol	m ²	cm ²
megabarn	Mb	10 ⁻²²	10 ⁻¹⁸
kilobarn	kb	10 ⁻²⁵	10 ⁻²¹
barn	b	10 ⁻²⁸	10 ⁻²⁴
millibarn	mb	10 ⁻³¹	10 ⁻²⁷
microbarn	μb	10 ⁻³⁴	10 ⁻³⁰
nanobarn	nb	10 ⁻³⁷	10 ⁻³³
picobarn	pb	10 ⁻⁴⁰	10 ⁻³⁶
femtobarn	fb	10 ⁻⁴³	10 ⁻³⁹
attobarn	ab	10 ⁻⁴⁶	10 ⁻⁴²
zeptobarn	zb	10 ⁻⁴⁹	10 ⁻⁴⁵
yoctobarn	yb	10 ⁻⁵²	10 ⁻⁴⁸



The Luminosity (~ Technology, not Physics)

$$\mathcal{L} = \phi_a \cdot N_b$$

Beam on a target

Luminosity : [(area x time)⁻¹]. From $\phi_a = n_a \times v_a$ and $N_b = n_b \cdot d \cdot A$ we have

$$\mathcal{L} = \phi_a \cdot N_b = \dot{N}_a \cdot n_b \cdot d = n_a \cdot v_a \cdot N_b$$

Luminosity → defined as one of two products below

1. number of incoming beam particles per unit time N_a , the target particle density in the scattering material n_b , and the target's thickness d ;
2. beam particle density n_a , their velocity v_a and the number of target particles N_b exposed to the beam.

j packets with N_a or N_b particles, a ring of circumference U . velocity $v \sim c$ in opposite directions and cross at an interaction point

$$\mathcal{L} = \frac{N_a \cdot N_b \cdot j \cdot v / U}{A}$$

two beams in a storage ring.

The luminosity is then:

A = beam cross-section at the collision point. For a Gaussian distribution of the beams (σ_x and σ_y respectively),

$$A = 4\pi\sigma_x\sigma_y.$$

... and have to be well aligned:
LHC ~27Km circumference!

→ beams must be focused at the interaction point into the smallest possible area possible. Typical beam diameters are of the order of tenths of millimetres or less.



Differential and Doubly-Differential Cross Sections

Real life: In all experiments only a fraction of all reactions are measured or accessible because of limited **acceptance** of the experimental set-up.


Detector of area A_D at a distance r and at an angle θ , it covers a solid angle equal to $\Delta\Omega = A_D/r^2$.

The reaction rate (assumed to depend on the energy of the incoming beam and on the angle θ) will be:

$$N(E, \theta, \Delta\Omega) = \mathcal{L} \frac{d\sigma(E, \vartheta)}{d\Omega} \Delta\Omega$$

If the energy & direction of the products is measured then the doubly differential cross section is also measured $d^2\sigma(E, E', \theta)/d\Omega dE'$. The total cross section, in this case, will be the integral over the solid angle and over the scattering energies

$$\sigma_{tot}(E) = \int_{E_{min}}^{E_{max}} \int_{\theta_{min}}^{\theta_{max}} \frac{d^2\sigma(E, E', \theta)}{d\Omega dE'} d\Omega dE'$$





Cross Sections from Theory: The Golden Rule

Scattering processes \rightarrow cross sections. Can we compute it with theory?

Particle Physics: based on the study of cross-sections and decays \rightarrow interaction rates & decay rates. These rates can be derived using the Fermi's Golden Rule.

Transition (or “reaction..”) rate from an initial ψ_i to a final state ψ_f

$$\Gamma_{fi} = 2\pi \cdot |M_{fi}|^2 \rho(E)$$

where M_{fi} (also T_{fi} in some cases) is

$$M_{fi} = \langle \psi_f | \mathcal{H}_{int} | \Psi_i \rangle + \text{higher orders}$$

And $\rho(E)$ is the ‘density of states’ (: “in how many possible ways can you create the final state you are studying?”)

- M_{fi} contains Physics
- $\rho(E)$ contains kinematics

If interaction rate is \sim low \rightarrow *perturbation expansion*



$$M_{fi} = \langle \psi_f | \mathcal{H}_{int} | \Psi_i \rangle$$



The Golden Rule: the Density of States

Cross section also depends on the number of final states available to the reaction.

According to the Heisenberg uncertainty principle each particle occupies a volume $h^3 = (2\pi\hbar)^3$ in the 6-dim position-momentum space $\Delta_x \cdot \Delta_y \cdot \Delta_z \cdot p_x \cdot p_y \cdot p_z$



Consider a particle scattered into a volume V in a momentum interval \mathbf{p}' , $\mathbf{p}'+\delta\mathbf{p}'$. The volume of this spherical shell is $4\pi p'^2 \delta p'$ and the total number of final states is

$$dn(p') = \frac{V 4\pi p'^2 \delta p'}{(2\pi\hbar)^3}$$

Volume occupied by 1 particle

the density of final states $\rho(E')$ is ($dE' = v' dp'$)

$$\rho(E') = \frac{dn}{dE'} = \frac{dn}{dp'} \frac{dp'}{dE'} = \frac{V 4\pi p'^2}{v' (2\pi\hbar)^3}$$

(you may consider $V = 1$)



The Golden Rule, the

According to the Fermi second golden rule (not derived here): *reaction rate* W (per beam particle and per target particle), transition matrix and density of final states

$$\Gamma_{fi} = 2\pi \cdot |M_{fi}|^2 \rho(E)$$

Few slides ago $\sigma_b = \frac{\dot{N}}{\phi_a \times N_b}$ ($\phi_a = n_a \times v_a$) $\rightarrow W = \frac{N(E)}{N_b N_a} = \frac{\sigma_b v_a}{V}$ where V = is the spatial volume occupied by beam particles

$$\sigma = \frac{2\pi}{\hbar v_a} \overset{\text{Theory}}{|M_{fi}|^2} \rho(E') V$$

- If interaction potential is known or calculable \rightarrow compute the cross section
- if M_{fi} is not known one can measure σ and derive M_{fi} from it.

The Golden Rule applies both to scattering and decay processes. In the second case the lifetime of the process will be

$$\tau = \frac{1}{W}$$

- if the lifetime is (can be) measured then M_{fi} can be derived.
- If τ cannot be measured then the uncertainty principle can be used and we can take $\Delta E = \hbar/\tau$







Feynman Diagrams (Povh...)

Feynman diagrams → graphic representation of scattering → symbols into the matrix element.

- Works for
 - QED and
 - weak interactions also for
 - strong interactions (quantum chromodynamics, QCD).

Drawing rules:



- The time axis runs \uparrow and the space axis \rightarrow .
- The straight lines correspond to the wave functions of the initial and final fermions.
- Antiparticles → arrows pointing backwards in time; by wavy lines;
- photons 
- heavy vector bosons by dashed lines; 
- and gluons by corkscrew-like lines. 


Uncertainty principle → virtual particles do not satisfy the energy-momentum relation $E^2 = p^2c^2 + m^2c^4$

- 1] the exchanged particle has a mass different from that of a free (real) particle, or
- 2] that energy conservation is violated for a brief period of time.




Feynman Diagrams for Em, Weak, Strong Interactions

Points at which three or more particles meet are called vertices.



Feynman diagrams for the electromagnetic (a, b, c), weak (d, e) and strong interactions (f).



In QED, as in other quantum field theories, we can use the little pictures invented by my colleague Richard Feynman, which are supposed to give the illusion of understanding what is going on in quantum field theory.

M. Gell-Mann [Ge80]




Feynman Diagrams – EM

$$\sigma = \frac{2\pi}{\hbar v_a} \hbar |M_{fi}|^2 \rho(E')V$$

$$\alpha = \frac{e^2}{\hbar c}$$

$$\sqrt{\alpha} \propto e$$



One vertex \rightarrow one term in the transition matrix (structure and strength of the interaction).


\leftarrow the exchanged photon couples to the charge of the electron at the left vertex and to that of the positron at the right vertex. For each vertex the transition amplitude contains a factor which is proportional to $e \rightarrow \sqrt{\alpha}$.

(b) \rightarrow annihilation of an e^+e^- pair. A photon is created as an intermediate state which then decays into a $\mu^+\mu^-$ pair.
 (c) same process with one additional diagram contributing to the same final state \rightarrow *higher-order diagrams*. The transition matrix is the sum of all diagrams \rightarrow same final state.

More vertices implies higher powers of α .

Diagram (b) is $\propto \alpha$ Diagram (c) is $\propto \alpha^2$.

$e^+e^- \rightarrow \mu^+\mu^-$ is given by graph (b) + higher diagrams produce only small corrections to (b). $\alpha \sim 1/137$






Feynman Diagrams – EM

Annihilation of an e^+e^- pair into a $\mu^+\mu^-$ pair.

=



=



Feynman Diagrams – Weak & Strong





Figure (d) shows electron-positron annihilation followed by muon pair production in a weak interaction: exchange of the neutral, heavy vector boson Z^0 → **weak interaction**



In Figure (e), we see a neutron that transform into a proton via β^- -decay in which it emits a negatively charged heavy vector boson W^- which then decays into an electron and antineutrino $\bar{\nu}_e$. → **weak interaction**

time ↑




Figure (f) strong interaction process between two quarks q and q' which exchange a gluon → **strong interaction**

(f)

(e)

(d)



Feynman Diagrams: Weak & Strong Interactions

- In weak interactions, a heavy vector boson is exchanged which couples to the “weak charge” g and not to the electric charge e . \rightarrow weak interaction, $M_{fi} \propto g^2 \propto \alpha_w$.
- In strong interactions the gluons which are exchanged between the quarks couple to the “colour charge” of the quarks, $M_{fi} \propto \sqrt{\alpha_s} \cdot \sqrt{\alpha_s} = \alpha_s$.

The ‘propagator’

The exchanged particles contribute a **propagator** term to the transition matrix element:

$$\frac{1}{Q^2 + M^2 c^2}$$

Q^2 is the four-momentum² of the virtual exchanged particle
 M is the mass of the virtual exchanged particle

Exchanged photon (EM interactions):

$\rightarrow 1/Q^2$ in the amplitude and $1/Q^4$ in the cross-section.

Exchanged W^\pm, Z (Weak interaction):

\rightarrow large mass \rightarrow the cross-section is much smaller than EM interaction

BUT at very high momentum transfers, of the order of the masses of the vector bosons, the two cross-sections become comparable in size.



Of the Scattering Processes (real life)


Scattering experiments → determine the transition matrix element.

Scattering experiment = beam projectiles particles hit a target (or two beams clashing against each other).


- **Projectiles are or may be extended objects** (low energy particles have a large $\lambda = \hbar/p$; The more energetic projectiles are used the smaller is the equivalent de Broglie wave length $\lambda = \hbar/p$, small wave lengths allow the inspection of the inner structure of the matter).
- protons have an **internal structure**, we know they are composite objects (alfa particles even more);

- **electrons**, as far as we know, **are point-like objects**, the interaction between electrons and a nucleus or a quark proceeds via the exchange of a photon;
- **Processes mediated by photons have two advantages:**
 1. The first one is that they are well known since long.
 2. The second positive fact is that these interactions are characterised by a strength (i.e. a coupling constant) $\alpha=1/137$ which is rather small and allows the perturbation

theory to be applied



higher level diagram → small contribution to cross section





Electron Nucleus Kinematics

Elastic Scattering of an Electron on a Particle at Rest with Mass M
(assumed to be a proton)

→ *electron and particle with mass M remain unchanged in the final state*


	Initial state	Final state
electron	E, p	E', p'
nucleus	E_p, P	E_p', P'

4 – Momenta !

The energy conservation implies that
 and also...

$$p + P = p' + P'$$

$$P' = p + P - p'$$



And, once squared,

$$\cancel{p^2} + 2pP + \cancel{P^2} = \cancel{p'^2} + 2p'P' + \cancel{P'^2}$$

4 momenta p^2 and p'^2 are invariant and equal to m_e
 4 momenta P^2 and P'^2 are also invariant and equal to M

$$p^2 = p'^2 = m_e^2 c^2 \quad \text{and} \quad P^2 = P'^2 = M^2 c^2$$

$$p \cdot P = p' \cdot (p + P - p') = p'p + p'P - m_e^2 c^2$$

Kinematics of elastic electron – nucleus scattering

$$p + P = p' + P' \rightarrow p' = p + P - P'$$




Electron Nucleus Kinematics

$$\mathbf{a} \cdot \mathbf{b} = a_0 b_0 - a_1 b_1 - a_2 b_2 - a_3 b_3 = a_0 b_0 - \mathbf{a} \cdot \mathbf{b}$$

only the electron is detected \rightarrow use electron kinematics

$$p \cdot P = p' \cdot (p + P - p') = p'p + p'P - m_e^2 c^2$$



Let's choose the laboratory frame where particle P is at rest \rightarrow

$$p = (E/c, \mathbf{p}) \quad p' = (E'/c, \mathbf{p}') \quad P = (Mc, \mathbf{0}) \quad P' = (E_P/c, \mathbf{P}')$$

$$E \cdot Mc^2 = E'E - \mathbf{p}\mathbf{p}'c^2 + E'Mc^2 - m_e^2 c^4$$

At high energy we may neglect the electron mass and take $E \sim |\mathbf{p}| \cdot c \rightarrow E \cdot Mc^2 = E'E \cdot (1 - \cos \theta) + E' \cdot Mc^2$

θ is the scattering angle between \mathbf{p} and \mathbf{p}'


$$E' = \frac{E}{1 + E/Mc^2 \cdot (1 - \cos \theta)}$$

In elastic scattering (and only there!) there is a one-to-one correlation between the scattering angle θ and the energy of the electron. The recoil energy transferred to the target proton is given by $E' - E$

\rightarrow if the term E/M increases E' decreases \rightarrow small recoil energy



Kinematics of Electron Scattering off Nucleus



$A = 50 \rightarrow \text{Sn (tin)}$

$$E' = \frac{E}{1 + E/Mc^2 \cdot (1 - \cos \theta)}$$

- Term $(1 - \cos \theta) E/Mc^2 \rightarrow$ angular dependence
- recoil energy increases with decreasing E/Mc^2

In electron scattering at 0.5 GeV off a nucleus with mass number $A = 50$ the scattering energy varies by only 2% at most.

At 10 GeV electrons scattering off protons ($A = 1$) E varies between 10 GeV ($\theta \approx 0^\circ$) and 445 MeV ($\theta = 180^\circ$).





Scattering of electrons on nucleus/proton

Calculation	electron		Target, charge Ze (Z=1 proton)					Expression (we visit these formulas in next slides)
	electron	Electron with spin	Point-like target, infinite Mass	Point-like target with mass M	Point-like proton	Point-like proton with spin	Finite size proton with spin	
Rutherford	✓		✓					$\left(\frac{d\sigma}{d\Omega}\right)_R = \frac{Z^2 e^4}{4E_0^2 (\sin \theta/2)^4}$
Mott		✓		✓				$\left(\frac{d\sigma}{d\Omega}\right)_M = \left(\frac{d\sigma}{d\Omega}\right)_R \cdot (\cos \frac{\theta}{2})^2$
σ_{NS}		✓			✓			$\left(\frac{d\sigma}{d\Omega}\right)_{NS} = \left(\frac{d\sigma}{d\Omega}\right)_M \cdot 1 / \left(1 - \frac{2E_0}{M} \sin \theta/2^2\right)$
σ		✓				✓		$\left(\frac{d\sigma}{d\Omega}\right) = \left(\frac{d\sigma}{d\Omega}\right)_M \cdot \left(1 + \frac{q^2}{2M^2} \tan \theta/2^2\right)$
Rosenbluth		✓					✓	$\left(\frac{d\sigma}{d\Omega}\right) = \left(\frac{d\sigma}{d\Omega}\right)_M \cdot \left[\frac{G_E^2(Q^2) + \tau \cdot G_M^2(Q^2)}{1 + \tau} + 2\tau G_M^2(Q^2) \tan \theta/2^2 \right]$


Account for, electron spin, target spin, charge distribution, magnetic moment

Expression
(we visit these formulas in next slides)





Rutherford scattering – Classical Calculation




Particles travelling at a large distance from the scattering centre are deflected less than particles travelling closer to the nucleus.

→ For a larger impact parameter b , the deflection occurs at a smaller angle. For this reason, a negative $d\theta$ corresponds to a positive db



Rutherford scattering – Classical Calculation



- Charged particles arrive on the ring ($2\pi b db$)
- Are elastic scattered by ~ a heavy nucleus which creates a Coulomb potential
- In the angular range $[\theta, \theta - \delta\theta]$

Compute this!

You measure this!

→ relation between the impact parameter b and the deflection angle in the Coulomb elastic scattering.

The number of incident particles elastically scattered per time unit in the interval $(\theta, \theta - \delta\theta)$ is

$$dN = N_0 2\pi b db = N_0 d\sigma \rightarrow d\sigma = 2\pi b db$$

$d\sigma(\theta)$ = cross section for scattering $(\theta \rightarrow \theta - \delta\theta)$

- N_0 is the number of incident particles per area and time units (*flux*)
- $d\sigma = 2\pi b db$ is the surface of the ring hit by the incident particles in the angular range $(\theta, \theta - \delta\theta)$.
- dN is number of particles arriving at b and scattered at θ



Rutherford Scattering - continued

Elastic differential cross-section:

$$d\sigma(\theta) = \frac{d\sigma}{d\Omega}(\theta) d\Omega = \frac{d\sigma}{d\Omega} \boxed{2\pi \sin(\theta)} d\theta$$

Integral over 2π in ϕ

$$\sigma_{tot}^{el} = \int \frac{d\sigma}{d\Omega}(\theta) d\Omega = 2\pi \int_0^\pi \frac{d\sigma}{d\Omega}(\theta) \sin(\theta) d\theta$$

Total cross-section: size of the “transverse” area of the diffusion centre

Scattering of few MeV α particles (${}^4\text{He}$ nuclei, $2p$, $\rightarrow z=2$) against gold ($Z = 79$) nuclei. The Coulomb potential nucleus (charge Z) + α particle (charge z) is

$$V(r) = zZe^2/r$$





Fig. 1.1 Schematic diagram of the apparatus used in the Rutherford scattering experiment. Alpha particles scattered by the gold foil strike a fluorescent screen, giving off a flash of light, which is observed visually through a microscope.



Rutherford scattering - continued



$$|\vec{p}'| = |\vec{p}|; \Delta p = |\vec{p}' - \vec{p}| = 2 p \sin\left(\frac{\theta}{2}\right)$$

After a bit of mathematics ... Look at Lecture Notes!!

p = momentum of α particle,

$$\tan\left(\frac{\theta}{2}\right) = \frac{zZe^2m}{p^2b} \rightarrow b = \frac{zZe^2 \cot(\theta/2)}{p^2} \frac{zZe^2m}{p^2 \tan(\theta/2)}$$

And using

- $d(\cot x) = -(\sin x)^2 dx$
- Introduce solid angle $d\Omega = 2\pi \sin\theta d\theta$
- and use $\sin\theta = 2 \sin\left(\frac{\theta}{2}\right) \cos\left(\frac{\theta}{2}\right)$
- $\frac{d\sigma}{d\Omega}(\theta) 2\pi \sin(\theta) d\theta = -2\pi b \cdot db$ (2 slides before)

$$\begin{aligned} \left(\frac{d\sigma}{d\Omega}\right)_{Rutherford} &= \\ \left(\frac{zZ^2e^2m}{4\pi\epsilon_0}\right)^2 \frac{1}{4p^4 \sin^4\left(\frac{\theta}{2}\right)} &= \\ = \left(\frac{zZ^2e^2m}{4\pi\epsilon_0}\right)^2 \frac{4}{(\Delta p)^4} \end{aligned}$$


Valid for a spinless particle





Rutherford scattering - continued

Look at Lecture Notes!!



$$|\vec{p}'| = |\vec{p}|; \Delta p = |\vec{p}' - \vec{p}| = 2 p \sin\left(\frac{\theta}{2}\right)$$

Classical Relation

$$\tan(\theta/2) = \frac{\text{Potential energy at a distance } 2b}{\text{Kinetic energy}}$$

$$= \frac{zZe^2}{2bE_c}$$

$$\rightarrow b = \frac{zZe^2 \cot(\theta/2)}{2E_c}$$

E_c = kinetic energy of α particle,

And using

- $d(\cot x) = -(\sin x)^2 dx$
- $\frac{d\sigma}{d\Omega}(\theta) 2\pi \sin(\theta) d\theta = -2\pi b \cdot db$ (2 slides before)

NEGLECT

$$\frac{db}{d\theta} = \frac{zZe^2}{2 \cdot 4 b E_c \sin(\theta/2)^2}$$

We get

$$\frac{d\sigma}{d\Omega}(\theta) = -\frac{b}{\sin\theta} \frac{db}{d\theta} = \frac{(zZe^2)^2}{(4 E_c)^2 \sin(\theta/2)^4}$$

← Valid for a spinless particle



Targets with Extended Charge Distribution

We treated a simple case, nucleus is point-like (\rightarrow low energy particles)

Let's consider the case of Extended Charge Distribution
 \rightarrow the probe sees the internal structure

The nucleus is NOT
point-like

- an electron beam with a density n_a particles per unit volume,
- scattering on a very heavy nucleus \rightarrow recoil is so small that ~ 0 .
- We use three-momenta.
- If target charge Ze is \sim small \rightarrow the electro-magnetic interaction will be small $Z\alpha \ll 1$ ($\alpha=1/137$).
- In this case the wave functions Ψ_i and Ψ_f of the initial and final state (i.e. electron) will be described by plane waves:

$$\Psi_i = \frac{1}{\sqrt{V}} e^{ipx/\hbar} \quad \Psi_f = \frac{1}{\sqrt{V}} e^{ip'x/\hbar}$$

We assume that the process takes place in a volume V (large with respect to the scattering centre) and that wave functions of the incoming and outgoing electrons are normalised in this volume. We have a total number of N_a electrons in the beam

$$\int_V |\psi_i|^2 dV = n_a \cdot V \quad \text{where} \quad V = \frac{N_a}{n_a}$$



Extended Charge Distribution

1) Scattering reaction rate:

$$W = \frac{\sigma v_a}{V}$$

2) We apply Fermi's Golden Rule:

$$W = \frac{2\pi}{\hbar} |M_{fi}|^2 \rho(E') = \frac{2\pi}{\hbar} |\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle|^2 \frac{dn}{dE_f}$$

3) Density of final states:

$$dn(p') = V 4\pi p'^2 dp' / (2\pi\hbar)^3$$

E_f is the total energy of the final state

If we combine 1, 2, 3

$$d\sigma \cdot v_a \cdot \frac{1}{V} = \frac{2\pi}{\hbar} |\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle|^2 \frac{V |\mathbf{p}'|^2 d|\mathbf{p}'|}{(2\pi\hbar)^3 dE_f} d\Omega$$

The beam particle velocity $v_a \sim c$ and $|\mathbf{p}'| \sim E'/c \rightarrow$

$$\frac{d\sigma}{d\Omega} = \frac{V^2 E'^2}{(2\pi)^2 (\hbar c)^4} |\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle|^2$$

Kinematics Physics



Some Calculation...

The interaction operator that transforms the initial state into a final one for a charge e in an electric potential ϕ is:

$$\mathcal{H}_{int} = e\phi(x)$$

And the matrix element $\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle$ becomes

$\phi \rightarrow \phi(x)$, diffused charge, not point-like

$$\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle = \frac{e}{V} \int e^{-\frac{i\mathbf{p}' \cdot \mathbf{x}}{\hbar}} \phi(x) e^{\frac{i\mathbf{p} \cdot \mathbf{x}}{\hbar}} d^3x$$

The momentum transfer between \mathbf{p} and \mathbf{p}' is defined as $\mathbf{q} = \mathbf{p} - \mathbf{p}' \rightarrow$

$$\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle = \frac{e}{V} \int \phi(x) e^{\frac{i\mathbf{q} \cdot \mathbf{x}}{\hbar}} d^3x$$

And with some calculation (Green's theorem & Poisson's equation, see next slide)

$$\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle = \frac{e\hbar^2}{\epsilon_0 V |\mathbf{q}^2|} \int f(x) e^{\frac{i\mathbf{q} \cdot \mathbf{x}}{\hbar}} d^3x = \frac{e\hbar^2}{\epsilon_0 V |\mathbf{q}^2|} F(\mathbf{q})$$

Where we have defined $F(\mathbf{q}) = \int f(x) e^{\frac{i\mathbf{q} \cdot \mathbf{x}}{\hbar}} d^3x$, called form factor of the charge distribution.

The form factor describes the charge distribution of the target we are studying in our scattering experiment.



Green's Theorem

■ Green's theorem permits us to use a clever trick here: for two arbitrarily chosen scalar fields u and v , which fall off fast enough at large distances, the following equation holds for a sufficiently large integration volume:

$$\int (u\Delta v - v\Delta u) d^3x = 0, \quad \text{with } \Delta = \nabla^2. \quad (5.27)$$

Inserting:

$$e^{i\mathbf{q}\mathbf{x}/\hbar} = \frac{-\hbar^2}{|\mathbf{q}|^2} \cdot \Delta e^{i\mathbf{q}\mathbf{x}/\hbar} \quad (5.28)$$


into (5.26), we may rewrite the matrix element as:

$$\langle \psi_f | \mathcal{H}_{\text{int}} | \psi_i \rangle = \frac{-e\hbar^2}{V|\mathbf{q}|^2} \int \Delta\phi(\mathbf{x}) e^{i\mathbf{q}\mathbf{x}/\hbar} d^3x. \quad (5.29)$$

The potential $\phi(\mathbf{x})$ and the charge density $\rho(\mathbf{x})$ are related by Poisson's equation:

$$\Delta\phi(\mathbf{x}) = \frac{-\rho(\mathbf{x})}{\epsilon_0}. \quad (5.30)$$

In the following, we will assume the charge density $\rho(\mathbf{x})$ to be static, i. e. independent of time.





Getting the point-like Rutherford Cross Section

$$\langle \psi_f | \mathcal{H}_{int} | \psi_i \rangle = \frac{e\hbar^2}{\epsilon_0 V |\mathbf{q}^2|} \int f(x) e^{\frac{i\mathbf{q}x}{\hbar}} d^3x = \frac{e\hbar^2}{\epsilon_0 V |\mathbf{q}^2|} F(\mathbf{q})$$

ϵ_0 is the permittivity of free space

If we neglect the fact that our target has an extended charge distribution then $F(\mathbf{q})$ becomes a δ function $\rightarrow F(\mathbf{q}) = 1$. If we do this approximation then we get Rutherford cross section in the case of a point-like charge distribution and expressed as a function of the momentum transfer \mathbf{q} .


$$d\sigma/d\Omega = \frac{4Z^2\alpha^2(\hbar c)^2 E'^2}{|\mathbf{q}c|^4}$$

$$\alpha_{EM} = \frac{e^2}{4\pi\epsilon_0\hbar c}$$

The $1/q^4$ dependence indicates that the cross section drops very quickly for large values of \mathbf{q} and that the largest part of the cross section is limited to small values of \mathbf{q} .

Let's remember that

- If we neglect recoil $E=E'$
- $|\mathbf{p}| = |\mathbf{p}'|$
- $E=|\mathbf{p}| c$ is a good approximation




$$|\mathbf{q}| = 2 \cdot |\mathbf{p}| \sin \frac{\theta}{2}$$

$$\left(\frac{d\sigma}{d\Omega} \right)_{\text{Rutherford}} = \frac{Z^2\alpha^2(\hbar c)^2}{4E^2 \sin^4 \frac{\theta}{2}}$$

We get the classical expression



Understanding $F(q^2)$



$q^2 \sim 0 \rightarrow$ point-like interaction, the nucleon is seen by the photon as a unique object $\rightarrow F(q^2) \sim 1$

$F(q^2)$ indicates how well the nucleon 'holds together' (= is not 'broken') if hit by a photon of momentum q^2

$q^2 \sim$ large \rightarrow diffused charge \rightarrow the photon sees only a part of the charge of the nucleon $\rightarrow F(q^2) \sim$ very low

When q^2 increases \rightarrow the probe becomes very small



Changing Point of View: Field Theory


The scattering of an electron on a target with charge Ze is mediated by the exchange of a virtual photon emitted by the electron and absorbed by the target (see picture here \rightarrow).

The transition matrix $\sim e \cdot Ze$ and the cross section $\sim (e \cdot Ze)^2$.

The momentum transfer q is the momentum carried by the photon and transferred from the electron to the target.

The equivalent de-Broglie wave-length of the photon is

$$\lambda = \frac{\hbar}{q} = \frac{\hbar}{|\mathbf{p}|} \frac{1}{2\sin\frac{\vartheta}{2}}$$







- *if the de-Broglie wave-length of the photon is NOT small enough with respect to the extension of the target it cannot probe the internal structure of the scattering centres and the target appears to be point-like.*
 - *The Rutherford cross section was obtained with low energy electrons corresponds to this situation.*

The propagator in the matrix element $\frac{1}{Q^2 + M^2 c^2}$ becomes simply $\frac{1}{Q^2}$



Rutherford Scattering





Scattering of electrons on nucleus/proton

Calculation	electron		Target, charge Ze (Z=1 proton)					Expression
	electron	Electron with spin	Point-like target, infinite Mass	Point-like target with mass M	Point-like proton	Point-like proton with spin	Finite size proton with spin	
Rutherford	✓		✓					$\left(\frac{d\sigma}{d\Omega}\right)_R = \frac{Z^2 e^4}{4E_0^2 (\sin \theta/2)^4}$
Mott		✓		✓				$\left(\frac{d\sigma}{d\Omega}\right)_M = \left(\frac{d\sigma}{d\Omega}\right)_R \cdot (\cos \frac{\theta}{2})^2$
σ_{NS}		✓			✓			$\left(\frac{d\sigma}{d\Omega}\right)_{NS} = \left(\frac{d\sigma}{d\Omega}\right)_M \cdot 1 / \left(1 - \frac{2E_0}{M} \sin \theta/2^2\right)$
σ		✓				✓		$\left(\frac{d\sigma}{d\Omega}\right) = \left(\frac{d\sigma}{d\Omega}\right)_M \cdot \left(1 + \frac{q^2}{2M^2} \tan \theta/2^2\right)$
Rosenbluth		✓					✓	$\left(\frac{d\sigma}{d\Omega}\right) = \left(\frac{d\sigma}{d\Omega}\right)_M \cdot \left[\frac{G_E^2(Q^2) + \tau \cdot G_M^2(Q^2)}{1 + \tau} + 2\tau G_M^2(Q^2) \tan \theta/2^2 \right]$





Electron Spin \rightarrow The Mott Cross Section (point-like objects)

The Rutherford cross section neglects the spin of the electron

The electron spin modifies the Rutherford cross section introducing a term $(1 - \beta^2 \sin^2 \frac{\theta}{2})$, $\beta = v/c$


$$(d\sigma/d\Omega)_{Mott} = (d\sigma/d\Omega)_{Rutherford} \cdot (1 - \beta^2 \sin^2 \frac{\theta}{2})$$

Large scattering angles are suppressed

In the limit case $\beta \rightarrow 1$ we get: $(d\sigma/d\Omega)_{Mott} = (d\sigma/d\Omega)_{Rutherford} \cdot (\cos^2 \frac{\theta}{2})$

\rightarrow for a scattering angle of 180° we have a zero cross-section

This is understood with the conservation of helicity, the projection of the spin in the direction of the motion : $h = \mathbf{s} \cdot \mathbf{p}/(|\mathbf{s}| \cdot |\mathbf{p}|)$



Neither the orbital angular momentum \mathbf{L} (pointing up with respect to the plane of motion) nor the spin-less target can compensate the flip of the helicity. The situation changes in case of a target with spin.



Nuclear Form Factors

- The Rutherford Cross Section represents well data only for small q .
- For higher values of q data are lower than predicted by formulas.
- This is understood with the fact that *assuming that the charge distribution $F(q)$ of a nucleus is point-like* is acceptable only when q is small \rightarrow the reduced wave length of the photon is too large to probe the charge distribution of the nucleus.
- In this picture (q small) the photon sees the nucleus (or the nucleon) as a unique object.
- When q increases the photon starts to see the inner structure of the proton \rightarrow the photon starts to see only a part of the charge and not all of it
- \rightarrow the cross section decreases with q faster than expected.

- The form factor, $F(q^2)$, carrying the information on how the charge is distributed inside the nucleus $f(x) \sim$ modulates the Mott cross section

$$F(q^2) = \int e^{\frac{iqx}{\hbar}} f(x) d^3x \quad \left(\frac{d\sigma}{d\Omega}\right)_{exp.} = \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \cdot |F(q^2)|^2$$

- The ratio between measurements and the Mott cross section for a point-like charge distribution allows the measurement of the charge distribution inside the nucleus.
- You measure the angle of the scattered electron, you compute q , you do the ratio.



More on Form Factors (Povh)

One could measure the ratio

$$|F(q^2)|^2 = \left(\frac{d\sigma}{d\Omega}\right)_{exp.} / \left(\frac{d\sigma}{d\Omega}\right)_{Mott}$$

But in practice one assumes different analytical shapes and compares data with predictions

Charge distribution $f(r)$		Form Factor $F(q^2)$	
point	$\delta(r)/4\pi$	1	constant
exponential	$(a^3/8\pi) \cdot \exp(-ar)$	$(1 + q^2/a^2\hbar^2)^{-2}$	dipole
Gaussian	$(a^2/2\pi)^{3/2} \cdot \exp(-a^2r^2/2)$	$\exp(-q^2/2a^2\hbar^2)$	Gaussian
homogeneous sphere	$\begin{cases} 3/4\pi R^3 & \text{for } r \leq R \\ 0 & \text{for } r > R \end{cases}$	$3\alpha^{-3}(\sin\alpha - \alpha\cos\alpha)$ with $\alpha = q R/\hbar$	oscillating





$$f(x) \quad F(q^2) = \int e^{\frac{iqx}{\hbar}} f(x) d^3x$$

		Example
pointlike	constant	Electron
	dipole	Proton
	gauss	${}^6\text{Li}$
homogeneous sphere	oscillating	-
sphere with a diffuse surface	oscillating	${}^{40}\text{Ca}$

$r \rightarrow$ $|q| \rightarrow$



Measuring the Charge Distribution of Nuclei





Magnetic Moments of Charged Spin $1/2$ Particles

Additional effect: interaction between the current of the electron and the nucleon's magnetic moment.

- The magnetic moment of a charged, spin $1/2$ point-like particle (a Dirac particle) is:

$$\mu = g \cdot \frac{e}{2M} \cdot \frac{\hbar}{2}$$

- M is the mass of the particle and the $g = 2$ factor is a result of relativistic quantum mechanics;
- The magnetic interaction is associated with a flip of the spin of the nucleon.
- Scattering through 0° is not allowed: conservation of both angular momentum and helicity and scattering through 180° is preferred. The magnetic interaction thus introduces into the interaction an **additional** factor containing a factor of $\tan^2 \theta/2$ and due to the interaction of the magnetic interaction proton/electron:

Interaction with charge of target

Interaction with magnetic moment of target

$$\left(\frac{d\sigma}{d\Omega}\right)_{point\ spin\ \frac{1}{2}} = \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \left[1 + 2\tau \tan^2 \frac{\theta}{2} \right]$$

$$\text{where } \tau = \frac{Q^2}{M^2 c^2}$$



Magnetic Moments of Charged Spin $\frac{1}{2}$ Particles

$$\left(\frac{d\sigma}{d\Omega}\right)_{point\ spin\ \frac{1}{2}} = \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \left[1 + 2\tau \tan^2 \frac{\theta}{2}\right] \quad \text{where } \tau = \frac{Q^2}{M^2 c^2}$$

- The interaction with magnetic moment of target is proportional to $1/(M^2)$, to the deflection of the electron (i.e., to the momentum transfer Q^2).
- The magnetic term in the expression is large at high four-momentum transfers Q^2 and if the scattering angle θ is large.

This additional term causes the cross section to fall off *less strongly at larger angles* \rightarrow at large values of Q^2 a flatter distribution is found than what predicted by the electric interaction.

The g-factor of a spin $\frac{1}{2}$ charged particle is exactly 2 (but for small understood very small deviations) while the g-factor of a spin $\frac{1}{2}$ neutral particle is exactly 0.

HOWEVER Nucleons are made up of quarks and this changes things
Their g-factors for protons and neutrons are determined by the internal structure

Measured!

$$\mu_p = \frac{g_p}{2} \mu_N = +2.79 \cdot \mu_N ,$$

$$\mu_n = \frac{g_n}{2} \mu_N = -1.91 \cdot \mu_N ,$$

Where μ_N is

$$\mu_N = \frac{e\hbar}{2M_p} = 3.1525 \cdot 10^{-14} \text{ MeV T}^{-1}$$



Magnetic Moments of the Proton and Neutron

Nuclei \leftrightarrow form factors ((nucleus has a structure))

Nucleons (with internal structure) \leftrightarrow form factors to describe electric $G_E(Q^2)$ and magnetic $G_M(Q^2)$ interactions (nucleon has a structure)


$$\left(\frac{d\sigma}{d\Omega}\right) = \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \cdot \left[\frac{G_E^2(Q^2) + \tau \cdot G_M^2(Q^2)}{1 + \tau} + 2\tau G_M^2(Q^2) \tan^2 \frac{\theta^2}{2} \right] \text{ where } \tau = \frac{Q^2}{M^2 c^2}$$

Q^2 dependence of the form factors \leftrightarrow the radial charge distributions and the magnetic moments.

- At low Q^2 $\left(\frac{d\sigma}{d\Omega}\right) / \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \approx G_E^2(Q^2)$
- At high Q^2 $\left(\frac{d\sigma}{d\Omega}\right) / \left(\frac{d\sigma}{d\Omega}\right)_{Mott} \approx \left(1 + 2\tau G_M^2(Q^2) \tan^2 \frac{\theta^2}{2}\right)$

\rightarrow Vary electron beam energy \rightarrow vary Q^2

\rightarrow measure electron scattering angle $\rightarrow \tan^2 \frac{\theta^2}{2}$





Introductory Part

Particle Physics
Toni Baroncelli
Haiping Peng
USTC

End of Introductory Part



Conservation Laws (~Invariance): Parity

- Invariance properties applies to physical systems described by an equation. The system is considered as invariant if the equation describing it is invariant under given transformations (say rotation or translation)
- Invariance properties are closely connected to conservation laws.

Transformations can be either continuous or discrete.

Symmetries are of great importance in physics. The conservation laws of classical physics (energy, momentum, angular momentum) are a consequence of the fact that the interactions are invariant with respect to their canonically conjugate quantities (time, space, angles). In other words, physical laws are independent of the time, the location and the orientation in space under which they take place.

Parity

Parity (P) is a reflection symmetry: depending on whether the sign of the wave function changes under reflection or not, the system is said to have negative or positive **P** respectively. For those laws of nature with left-right symmetry, the parity quantum number **P** of the system is conserved.

inverts the coordinates

does not change time

as a consequence

it inverts momenta

and does not change angular momenta

including spins

$$\mathbf{r} \Rightarrow -\mathbf{r}$$

$$t \Rightarrow t$$

$$\mathbf{p} \Rightarrow -\mathbf{p}$$

$$\mathbf{r} \times \mathbf{p} \Rightarrow \mathbf{r} \times \mathbf{p}$$

$$\mathbf{s} \Rightarrow \mathbf{s}.$$



Symmetries in Particle Physics: Parity

One has to ascribe an intrinsic parity \mathbb{P} to particles and antiparticles.

- Bosons and anti-bosons have the same intrinsic parity.
- fermions and antifermions have opposite parities.
- For a many-body system, **\mathbb{P} is a multiplicative quantum number** :


$$\mathbb{P}\psi(\vec{x}_1, \vec{x}_2 \dots \vec{x}_n, t) = P_1 P_2 \dots P_n \psi(\vec{x}_1, \vec{x}_2 \dots \vec{x}_n, t).$$

- Particles in a state of orbital angular momentum are parity eigenstates :


$$Y_{km}(\theta, \phi) = (-1)^k Y_{km}(\pi - \theta, \phi + \pi) \rightarrow \mathbb{P} |\psi_{km}(\theta, \phi)\rangle = (-1)^k |\psi_{km}(\theta, \phi)\rangle$$

- Therefore, for a two or a three particle system:

$$P_{\text{sys}(12)} = P_1 P_2 (-1)^L ;$$



$$P_{\text{sys}(123)} = P_1 P_2 P_3 (-1)^{L_1 + L_2}$$



It is an experimental fact that parity is conserved in all transitions due to electromagnetic and strong interactions. Parity is instead violated in weak interaction transitions



Wave function for two identical particles

The concept of parity has been generalised in relativistic quantum mechanics. One has to ascribe an intrinsic parity \mathbf{P} to particles and antiparticles.

Take a system with *two identical particles* and define an operator I that exchanges the two particles

$$I(1,2) \rightarrow (2,1)$$

The corresponding wave function will not change

$$I\Psi(1,2) = \Psi(2,1)$$

If we apply “I” twice we go back to the initial situation

$$I^2\Psi(1,2) = \Psi(1,2) \rightarrow \text{possible eigenvalues of } I \text{ are } \pm 1$$

It is assumed that


- Bosons and antibosons have integer spin (follow the Bose-Einstein statistics) and the same intrinsic parity. $\Psi(1,2) = \Psi(2,1)$ **are symmetric**
- Fermions and antifermions have half integer spin and opposite parities. $\Psi(1,2) = -\Psi(2,1)$ **anti-symmetric**

- Convention: P(quarks/leptons) =
+1 = $P_{e^-} = P_{\mu^-} = P_{\tau^-} = P_u = P_d = P_s = \dots$;
-1 = $P_{e^+} = P_{\mu^+} = P_{\tau^+} = P_{\bar{u}} = P_{\bar{d}} = P_{\bar{s}} = \dots$



Getting the Breit Wigner Shape

Let us imagine the elastic formation process of a generic resonance R, which decays with lifetime τ into the same initial particles. The presence of a interaction process is demonstrated by the different directions and momenta of the particles in the final state, that is,



The unstable resonance R is described by the free particle wave function $\psi(0)e^{-i\omega_R t}$ multiplied by a real function describing its decay probability as a function of time, that is,

$$\psi(t) = \boxed{\psi(0)e^{-i\omega_R t}} \boxed{e^{-\frac{t}{\tau}}} = \psi(0)e^{-\frac{iE_R}{\hbar}t} e^{-\frac{\Gamma}{2\hbar}t},$$

where the relations $\omega_R = \frac{E_R}{\hbar}$ and $\tau = \frac{\hbar}{\Gamma}$ have been inserted in the last equality. The probability of finding the particle at a time t is

$$I(t) = \psi^* \psi = \psi(0)^2 e^{-t/\tau} = I(0) \boxed{e^{-t/\tau}}, \quad \text{Exponential life-time}$$

corresponding to the radioactive decay law.

The Fourier transform is a transformation that decomposes functions depending on space or time into functions depending on spatial or temporal frequency \rightarrow gives us the energy distribution

$$\chi(E) = \int \psi(t)e^{iEt} dt = \psi(0) \int e^{-t[(\Gamma/2)+iE_R-iE]} dt = \int_0^\infty e^{-ax} dx = \frac{1}{a} = \frac{\boxed{K \ ?}}{(E_R - E) - i\Gamma/2}$$



The Transition Matrix Element

The cross-section can be experimentally determined from the reaction rate \dot{N} , as we saw above. We now outline how it may be found from theory. First, the reaction rate is dependent upon the properties of the interaction potential described by the Hamilton operator \mathcal{H}_{int} . In a reaction, this potential transforms the initial-state wave function ψ_i into the final-state wave function ψ_f . The transition matrix element is given by:

$$\mathcal{M}_{fi} = \langle \psi_f | \mathcal{H}_{\text{int}} | \psi_i \rangle = \int \psi_f^* \mathcal{H}_{\text{int}} \psi_i \, dV .$$

Furthermore, the reaction rate will depend upon the number of final states available to the reaction. According to the uncertainty principle, each particle occupies a volume $h^3 = (2\pi)^3$ in phase space, the six-dimensional space of momentum and position. Consider a particle scattered into a volume V and into a momentum interval between p and $p + dp$. In momentum space, the interval corresponds to a spherical shell with inner radius p and thickness dp which has a volume $4\pi p^2 dp$. Excluding processes where the spin changes, the number of final states available is:

$$dn(p') = \frac{V \cdot 4\pi p'^2}{(2\pi\hbar)^3} dp' .$$



The Transition Matrix Element

The energy and momentum of a particle are connected by:

$$dE' = v' dp'.$$

Hence the density of final states in the energy interval dE is given by:

$$\rho(E') = \frac{dn(E')}{dE'} = \frac{V \cdot 4\pi p'^2}{v' \cdot (2\pi\hbar)^3}.$$

The connection between the reaction rate, the transition matrix element and the density of final states is expressed by Fermi's second golden rule (not discussed here). It expresses the reaction rate W per target particle and per beam particle in the form:

$$W = \frac{2\pi}{\hbar} |\mathcal{M}_{fi}|^2 \cdot \rho(E').$$

with cross section

$$\sigma = \frac{2\pi}{\hbar \cdot v_a} |\mathcal{M}_{fi}|^2 \cdot \rho(E') \cdot V.$$



The Transition Matrix Element

The golden rule applies to both scattering, to the decay of unstable particles, to the excitation of particle resonances and to the transitions between different atomic or nuclear energy states. In these cases we have

$$W = \frac{1}{\tau}$$

and the transition probability per unit time can be either directly determined by measuring the lifetime τ or indirectly read off from the energy width of the state

$$\Delta E = \hbar/\tau$$