



ALICE

Measurements of inclusive $\psi(2S)$ to J/ψ ratio at midrapidity in pp collisions at $\sqrt{s} = 13.6$ TeV with ALICE

Yuan Zhang

University of Science and Technology of China

yuan.z@cern.ch




Abstract


Quarkonium production in high-energy pp collisions is an important tool for studying perturbative and non-perturbative aspects of quantum chromodynamics (QCD) calculations. The production process of charmonia can be factorized into two stages: the heavy quark production and the formation of the bound state. The former happens within initial hard parton-parton scatterings with large momentum transfers, and can be well described by perturbative QCD. The second one, which involves long distances and soft momentum scales, is a typical non-perturbative process. Measurements of J/ψ and $\psi(2S)$ cross section in pp collisions are crucial for studying charmonium production mechanisms and testing different QCD-based model calculations. They can also provide a reference for investigating the quark-gluon plasma formed in nucleus-nucleus collisions and the cold nuclear matter effects present in proton-nucleus collisions.

In this poster, the ratio of production of $\psi(2S)$ and J/ψ is shown based on the data collected in 2022 by the upgraded ALICE detector during the Run 3 of LHC, which offers significantly higher statistics compared to previous runs. The result is compared with measurements from other experiments at different energy and also several model calculations.

Motivation




- Crucial for studying charmonium production mechanisms and testing different QCD-based models.
 - Heavy-quark production (perturbative QCD)
 - Formation of the bound charmonium states (non-perturbative QCD)




- Study the rapidity dependence of charmonium production by comparing to similar measurements at forward rapidity at the same collision energy^[1].
- Used as reference for studying AA collisions.

ALICE detector (Run 2 configuration and Run 3 upgrade)

- Inclusive quarkonia are reconstructed in e^+e^- channel at midrapidity ($|y| < 0.9$) down to $p_T = 0$.



Data analysis procedure



- Dataset:
 - pp collisions at $\sqrt{s} = 13.6$ TeV collected in 2022 with the ALICE upgraded detector.
 - 524×10^9 minimum-bias (MB) events collected thanks to the continuous readout.
- Electron identification via TPC dE/dx .
- Signal extraction:
 - Combinatorial background is subtracted using mixed-event unlike-sign method.
 - Residual background: second order polynomial function divided by an exponential function.
 - Signal shapes are described by Double Crystal Ball functions. Possible differences between the J/ψ and $\psi(2S)$ shapes are assigned as systematic uncertainties.



- Efficiency correction:
 - Tracking efficiency and efficiency related to the choice of the signal mass window largely cancel out in the $\psi(2S)$ -to- J/ψ ratio. Residuals are assigned as systematic uncertainties.
 - PID efficiency is assessed using a data-driven approach.
 - Acceptance effects are corrected with a MC simulation.

Results



$$\frac{\sigma_{\psi(2S)}}{\sigma_{J/\psi}} = \frac{N_{\psi(2S)}}{N_{J/\psi}} \frac{(A \times \epsilon)_{J/\psi}}{(A \times \epsilon)_{\psi(2S)}} \frac{BR_{J/\psi \rightarrow ee}}{BR_{\psi(2S) \rightarrow ee}}$$

- N: raw counts.
- $A \times \epsilon$: acceptance times efficiency
- BR: Branching ratio

- The measured p_T -integrated ratio without BR uncertainty is $0.155 \pm 0.010(\text{stat.}) \pm 0.014(\text{syst.})$



- The results (red points) are shown together with existing results from ALICE at forward rapidity and from other experiments^[1-10].
- In agreement with other results.
- No significant energy and rapidity dependence.
- Slight p_T dependence (also expected from models).



- Comparison with models^[11-14]:
 - NRQCD overestimates the ratio at high p_T .
 - CGC + NRQCD describes the ratio at low and intermediate p_T .
 - ICEM can reproduce the data.

Summary and outlook

- The $\psi(2S)$ -to- J/ψ ratio is measured in pp collision at $\sqrt{s} = 13.6$ TeV at midrapidity.
 - In agreement with other results.
 - A slight p_T dependence (also expected from models).
 - No significant energy and rapidity dependence.
 - Comparison with models^[11-14].
 - NRQCD overestimates the ratio.
 - CGC + NRQCD describes the ratio at low and intermediate p_T .
 - ICEM can reproduce the data.
- Provides a reference for investigating the quark-gluon plasma in nucleus-nucleus collisions and the cold nuclear matter effects in proton-nucleus collisions.
- The prompt and non-prompt $\psi(2S)$ -to- J/ψ ratio as well as the cross section of prompt/non-prompt charmonia will be measured in Run 3.

Reference

- [1] ALICE Collaboration, S. Acharya et al., Eur. Phys. J. C 83 (2023) 61
- [2] LHCb Collaboration, R. Aaij et al., J. Phys. G 40 (2013) 045001.
- [3] E705 Collaboration, L. Antoniazzi et al., Phys.Rev.Lett. 70 (1993) 383-386.
- [4] STAR Collaboration, J. Adam et al., Phys. Rev. D 100 (2019) 052009.
- [5] PHENIX Collaboration, A. Adare et al., Phys. Rev. D 85 (2012) 092004.
- [6] A.G. Clark et al., Nucl. Phys. B 142 (1978) 29.
- [7] HERA-B Collaboration, I. Abt et al., Eur.Phys.J.C 49 (2007) 545-558.
- [8] CMS Collaboration, S. Chatrchyan et al., JHEP 02 (2012) 011.
- [9] CDF Collaboration, F. Abe et al., Phys.Rev.Lett. 79 (1997) 572-577.
- [10] ATLAS Collaboration, G. Aad et al., Eur.Phys.J.C 76 (2016) 5, 283.
- [11] Y.-Q. Ma et al., Phys.Rev.Lett. 106 (2011) 042002.
- [12] M. Butenschön et al., Phys.Rev.Lett. 106 (2011) 022003.
- [13] Y.-Q. Ma et al., Phys.Rev.D 94 (2016) 11, 114029.
- [14] M. Cacciari et al., JHEP 10 (2012) 137.

QPT 2023

