

Sensitivity for parameters of semileptonic hyperon decays

Varvara Batozskaya^{1,2}, Andrzej Kupść^{2,3}

¹Institute of High Energy Physics, Beijing, China

²National Centre for Nuclear Research, Warsaw, Poland

³Uppsala University, Uppsala, Sweden

Semileptonic meeting

13 May 2021

Sensitivity for extracted parameters

- Study the importance of the individual parameters in joint angular distribution and its correlations using likelihood function [\[PRDD100\(2019\)11\]](#)

$$\mathcal{L}(\omega) = \prod_{i=1}^N \mathcal{P}(\xi_i, \omega) \equiv \prod_{i=1}^N \frac{\mathcal{W}(\xi_i, \omega)}{\int \mathcal{W}(\xi, \omega) d\xi}$$

N is the number of events in the final selection

ξ_i is the full set of kinematic variables describing i -th event

ω is the full set of individual parameters

- Reduced asymptotic expression of inverse covariant matrix element:

$$V_{kl}^{-1} = N \int \frac{1}{\mathcal{P}} \frac{\partial \mathcal{P}}{\partial \omega_k} \frac{\partial \mathcal{P}}{\partial \omega_l} d\xi$$

- **Sensitivity** = $\sigma \times \sqrt{N}$
- To eliminate complicated calculation of derivatives the numerical differentiation is used [\[Numerical Analysis\]](#):

$$\frac{\partial \mathcal{P}}{\partial \omega_k} = \sum_{\omega_1}^{\omega_n} \lim_{h \rightarrow 0} \frac{\mathcal{P}(\omega_k + h) - \mathcal{P}(\omega_k)}{h}$$

Definitions and input values

$$A_D \equiv \frac{X_D + X_{\bar{D}}}{X_D - X_{\bar{D}}} \quad \text{and} \quad \langle X_D \rangle \equiv \frac{X_D - X_{\bar{D}}}{2}$$

where $X_D = \alpha_\Lambda, g_{av}^\Lambda, g_w^\Lambda$ and $X_{\bar{D}} = \alpha_{\bar{\Lambda}}, g_{av}^{\bar{\Lambda}}, g_w^{\bar{\Lambda}}$

- Input values of decay parameters [[NaturePhys.15\(2019\)631](#)] and slide 19

Decay	α_ψ	$\Delta\Phi$	α_Λ	g_{av}	g_w
$J/\psi \rightarrow \Lambda\bar{\Lambda}$	$0.461 \pm 0.006 \pm 0.007$	$0.740 \pm 0.010 \pm 0.008$	0.750 ± 0.010	0.719	1.066
$\Lambda \rightarrow p\pi^-$					
$\Lambda \rightarrow pe^-\bar{\nu}_e$					

- Decay parameters with the same value and opposite sign are used for $\bar{\Lambda}$ decays

Sensitivity ($\sigma \times \sqrt{N}$) for $\Lambda \rightarrow p e \nu_e$ decay

- Sensitivity for $\Lambda \bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\pi^+)$ is in an agreement with [\[PRD100\(2019\)114005\]](#)

Decay	α_ψ	$\Delta\Phi$	α_Λ	g_{av}^Λ	g_w^Λ	$\langle\alpha_\Lambda\rangle$	A_Λ	$\langle g_{av}^\Lambda\rangle$	A_{av}^Λ	$\langle g_w^\Lambda\rangle$	A_w^Λ
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\pi^+)$	3.43	7.47	6.83			1.76	8.81				
$\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}\pi^+)$ ($q^2 = 0, g_w = 0$)	3.19	7.13	6.97	21.0							
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}e^+\nu_e)$ ($q^2 = 0, g_w = 0$)	3.19	7.13	6.97	20.9							
$\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}\pi^+)$ ($g_w = 0$)	3.46	7.57	3.46	10.6							
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}e^+\nu_e)$ ($g_w = 0$)	3.46	7.57	3.46	10.6							
$\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}\pi^+)$	3.50	7.74	3.73	25.9	120						
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}e^+\nu_e)$	3.40	7.36	3.55	21.0	115						
combination ($q^2 = 0, g_w = 0$)						4.93	6.57	14.8	20.6		
combination ($g_w = 0$)						2.45	3.26	7.47	10.4		
combination						2.56	3.43	16.7	23.2	83.2	78.1
$\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}e^+\nu_e)$ ($q^2 = 0, g_w = 0$)	2.79	6.61		21.1				2.98	28.9		
$\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}e^+\nu_e)$ ($g_w = 0$)	3.42	7.42		10.6				7.33	10.6		
$\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}e^+\nu_e)$	3.43	7.44		24.2	111			15.6	22.9	83.5	69.5

- Correlations between parameters

$(q^2 = 0, g_w = 0)$						$(g_w = 0)$												
	g_{av}^Λ	g_{av}^Λ	$\alpha_{\bar{\Lambda}}$	α_ψ	$\Delta\Phi$		g_{av}^Λ	g_{av}^Λ	$\alpha_{\bar{\Lambda}}$	α_ψ	$\Delta\Phi$	g_{av}^Λ	g_{av}^Λ	g_{av}^Λ	g_w^Λ	$\alpha_{\bar{\Lambda}}$	α_ψ	$\Delta\Phi$
g_{av}^Λ	1	0.96		-0.02	0.02	g_{av}^Λ	1	0.04		0.04	0.05	g_{av}^Λ	1	0.05	-0.86	0.08	-0.04	-0.04
g_{av}^Λ		1		0.05	0.03	g_{av}^Λ		1		-0.04	-0.05	g_{av}^Λ		1	-0.08	0.91	0.04	0.05
$\alpha_{\bar{\Lambda}}$	0.93		1			$\alpha_{\bar{\Lambda}}$	-0.18		1			g_w^Λ	-0.87		1	-0.12	0.06	0.07
α_ψ	-0.06		0.01	1	0.02	α_ψ	0.01	0.19	1	0.24		$\alpha_{\bar{\Lambda}}$	0.21	-0.31		1		
$\Delta\Phi$	0.04		0.10	0.18	1	$\Delta\Phi$	0.01	0.25	0.27	1		α_ψ	-0.01	0.01		0.17	1	0.25
												$\Delta\Phi$	-0.00	0.00		0.24	0.28	1

* $\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}e^+\nu_e)$ (above the diagonal) and $\Lambda\bar{\Lambda} \rightarrow (pe^-\bar{\nu}_e)(\bar{p}\pi^+)$ (below the diagonal)

Sensitivity ($\sigma \times \sqrt{N}$) for $\Lambda \rightarrow p\mu\nu_\mu$ decay (1)

- Sensitivity for $\Lambda \rightarrow p e \nu_e$ and $\Lambda \rightarrow p \mu \nu_\mu$ is the same for $g_w^\Lambda = q^2 = 0$

Decay	α_ψ	$\Delta\Phi$	α_Λ	g_{av}^Λ	g_w^Λ	$\langle\alpha_\Lambda\rangle$	A_Λ	$\langle g_{av}^\Lambda \rangle$	A_{av}^Λ	$\langle g_w^\Lambda \rangle$	A_w^Λ
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\pi^+)(g_w = 0)$	3.51	7.77	3.72	10.5							
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\mu^+\nu_\mu)(g_w = 0)$	3.51	7.77	3.71	10.4							
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\pi^+)$	3.56	7.95	3.84	41.9	192						
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\mu^+\nu_\mu)$	3.46	7.56	3.60	29.8	165						
combination ($g_w = 0$)						2.63	3.50	7.39	10.3		
combination						2.63	3.51	25.7	35.7	126	119
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\mu^+\nu_\mu)(g_w = 0)$	3.54	7.83		10.6				7.01	10.9		
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\mu^+\nu_\mu)$	3.54	7.80		41.3	191			25.8	36.5	130	122

- Correlations between parameters

$(g_w = 0)$							g_{av}^Λ	g_{av}^Λ	g_w^Λ	g_w^Λ	$\alpha_{\bar{\Lambda}}$	α_ψ	$\Delta\Phi$
	g_{av}^Λ	g_{av}^Λ	$\alpha_{\bar{\Lambda}}$	α_ψ	$\Delta\Phi$	g_{av}^Λ	1	0.02	-0.95	—		0.03	0.04
g_{av}^Λ	1	0.11		0.06	0.09	g_{av}^Λ		1	-0.01	0.96		-0.02	-0.03
g_{av}^Λ		1		-0.06	-0.09	g_w^Λ	-0.95		1	—		-0.02	-0.02
$\alpha_{\bar{\Lambda}}$	-0.31		1			g_w^Λ				1		—	—
α_ψ	0.01		0.17	1	0.25	$\alpha_{\bar{\Lambda}}$	-0.18		0.10		1		
$\Delta\Phi$	0.01		0.25	0.28	1	α_ψ	0.01		-0.00		0.17	1	0.25
						$\Delta\Phi$	-0.00		0.00		0.26	0.28	1

* $\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\mu^+\nu_\mu)$ (above the diagonal) and $\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\pi^+)$ (below the diagonal)

Sensitivity ($\sigma \times \sqrt{N}$) for $\Lambda \rightarrow p\mu\nu_\mu$ decay (2)

- Additional factor $p = \sqrt{Q_+ Q_-} / (2M_1) \rightarrow p \frac{(q^2 - m_\mu^2)^2}{q^2}$
where $Q_\pm = (M_1 \pm M_2)^2 - q^2$

Decay	α_ψ	$\Delta\Phi$	α_Λ	g_{av}^Λ	g_w^Λ	$\langle\alpha_\Lambda\rangle$	A_Λ	$\langle g_{av}^\Lambda \rangle$	A_{av}^Λ	$\langle g_w^\Lambda \rangle$	A_w^Λ
Factor: p											
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\pi^+)(g_w = 0)$	3.51	7.77	3.72	10.5							
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\mu^+\nu_\mu)(g_w = 0)$	3.51	7.77	3.71	10.4							
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\pi^+)$	3.56	7.95	3.84	41.9	192						
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\mu^+\nu_\mu)$	3.46	7.56	3.60	29.8	165						
combination ($g_w = 0$)						2.63	3.50	7.39	10.3		
combination						2.63	3.51	25.7	35.7	126	119
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\mu^+\nu_\mu)(g_w = 0)$	3.54	7.83		10.6				7.01	10.9		
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\mu^+\nu_\mu)$	3.54	7.80		41.3	191			25.8	36.5	130	122
Factor: $p \frac{(q^2 - m_\mu^2)^2}{q^2}$											
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\pi^+)(g_w = 0)$	3.57	8.02	3.92	10.2							
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\mu^+\nu_\mu)(g_w = 0)$	3.57	8.01	3.92	10.2							
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\pi^+)$	3.61	8.17	4.41	52.7	268						
$\Lambda\bar{\Lambda} \rightarrow (p\pi^-)(\bar{p}\mu^+\nu_\mu)$	3.53	7.83	3.96	35.6	221						
combination ($g_w = 0$)						2.77	3.70	7.18	9.99		
combination						2.96	3.95	31.8	44.2	174	163
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\mu^+\nu_\mu)(g_w = 0)$	3.67	8.30		10.6				6.95	11.0		
$\Lambda\bar{\Lambda} \rightarrow (p\mu^-\bar{\nu}_\mu)(\bar{p}\mu^+\nu_\mu)$	3.67	8.33		50.2	262			28.8	47.3	188	157

Sensitivity ($\sigma \times \sqrt{N}$) for $\Lambda \rightarrow p\mu\nu_\mu$ decay (3)

- Correlations are the same for $g_w^\Lambda = 0$ and both factors
- Correlations between parameters for $g_w^\Lambda \neq 0$ and full q^2 range

factor p								factor $p \frac{(q^2 - m_\mu^2)^2}{q^2}$							
	g_{av}^Λ	g_{av}^Λ	g_w^Λ	g_w^Λ	$\alpha_{\bar{\Lambda}}$	α_ψ	$\Delta\Phi$		g_{av}^Λ	g_{av}^Λ	g_w^Λ	g_w^Λ	$\alpha_{\bar{\Lambda}}$	α_ψ	$\Delta\Phi$
g_{av}^Λ	1	0.02	-0.95	—		0.03	0.04	g_{av}^Λ	1	0.17	-0.97	0.13		0.08	0.11
g_{av}^Λ		1	-0.01	0.96		-0.02	-0.03	g_{av}^Λ		1	-0.15	0.97		-0.05	-0.08
g_w^Λ	-0.95		1	—		-0.02	-0.02	g_w^Λ	-0.97		1	-0.11		-0.07	-0.09
g_w^Λ				1		—	—	g_w^Λ				1		-0.03	-0.06
$\alpha_{\bar{\Lambda}}$	-0.18		0.10		1			$\alpha_{\bar{\Lambda}}$	-0.47		0.41		1		
α_ψ	0.01		-0.00		0.17	1	0.25	α_ψ	0.02		-0.02		0.14	1	0.26
$\Delta\Phi$	-0.00		0.00		0.26	0.28	1	$\Delta\Phi$	-0.01		0.01		0.25	0.28	1

* $\Lambda\bar{\Lambda} \rightarrow (p\mu^- \bar{\nu}_\mu)(\bar{p}\mu^+ \nu_\mu)$ (above the diagonal) and $\Lambda\bar{\Lambda} \rightarrow (p\mu^- \bar{\nu}_\mu)(\bar{p}\pi^+)$ (below the diagonal)

MC generator

- Use `YYbar_example` package by P.Adlarson to generate:

- $e^+e^- \rightarrow (\Lambda \rightarrow p e^- \bar{\nu}_e)(\bar{\Lambda} \rightarrow \bar{p} \pi^+)$ process
- $e^+e^- \rightarrow (\Lambda \rightarrow p \mu^- \bar{\nu}_\mu)(\bar{\Lambda} \rightarrow \bar{p} \pi^+)$ process:
 - factor: p
 - factor: $p \frac{(q^2 - m_\mu^2)^2}{q^2}$
- MC samples: $N_{\text{evt}}^{\text{sig}} = 10^5$ and $N_{\text{evt}}^{\text{phsp}} = 10^6$
- Generate $q^2 \in (m_l^2, (M_\Lambda - M_p)^2)$
- Set of input values:

```
pp[0] = 0.461;      // alpha_J/psi (arxiv:1808.08917)
pp[1] = 0.74;       // Delta_Phi (arxiv:1808.08917, equal 42.4deg)
pp[2] = 0.719;      // gav Lambda->p e- nu_ebar
pp[3] = 1.066;      // gw Lambda->p e- nu_ebar
pp[4] = -0.758;     // alpha_D LambdaBar->pbar pi+ (arxiv:1808.08917)
```

- No negative weights are observed
- Maximal weight ~ 0.36 for e -mode, ~ 0.3 and ~ 0.056 for μ -mode for two factors, respectively

MLL fit results

- Output value of fit method

Parameter	$N_{sig}^{MC} = 10^5, N_{phsp}^{MC} = 10^6$		
	e -mode	μ -mode (p)	μ -mode ($p \frac{(q^2 - m_\mu^2)^2}{q^2}$)
α_Ψ	0.4570 ± 0.0110	0.4563 ± 0.0120	0.4397 ± 0.0112
$\Delta\Phi$	0.7927 ± 0.0252	0.8232 ± 0.0267	0.7303 ± 0.0249
g_{av}	0.6601 ± 0.0506	0.6543 ± 0.0656	0.7294 ± 0.0870
g_w	1.1251 ± 0.2727	1.2275 ± 0.3306	1.2399 ± 0.4844
$\alpha_{\bar{\Lambda}}$	-0.7527 ± 0.0121	-0.7365 ± 0.0118	-0.7739 ± 0.0126

$\Lambda \rightarrow p e \nu_e$						$\Lambda \rightarrow p \mu \nu_\mu$					
	α_Ψ	$\Delta\Phi$	g_{av}	g_w	$\alpha_{\bar{\Lambda}}$		α_Ψ	$\Delta\Phi$	g_{av}	g_w	$\alpha_{\bar{\Lambda}}$
α_Ψ	1	0.281	-0.026	0.032	0.151	α_Ψ	1	0.288	-0.014	0.026	0.155
$\Delta\Phi$		1	0.014	-0.005	0.290	$\Delta\Phi$	0.266	1	0.016	-0.013	0.269
g_{av}			1	-0.848	0.096	g_{av}	0.016	-0.010	1	-0.766	0.323
g_w				1	-0.206	g_w	-0.016	0.006	-0.914	1	-0.411
$\alpha_{\bar{\Lambda}}$					1	$\alpha_{\bar{\Lambda}}$	0.135	0.274	-0.269	0.162	1

- Similar result for both modes and both factors

ToDo list and next steps


- Test formalism using
 - 1 Production of **mDIY** and **MC PhSp** for $e^+e^- \rightarrow (\Lambda \rightarrow p\pi^-)(\bar{\Lambda} \rightarrow \bar{p}\pi^+)$
 - true and reco **mDIY** and **MC PhSp**
 - allow to extract true and reco values of α_Λ and $\alpha_{\bar{\Lambda}}$ decay parameters
 - ♠ $\alpha_\Lambda^{\text{true}}$ and $\alpha_{\bar{\Lambda}}^{\text{true}}$ are extracted and verified
 - 2 Modification of **mDIY** to include the semileptonic decay formalism
 - ♠ Done
 - 3 Production of true and reco **mDIY** and **MC PhSp** for $\Lambda \rightarrow pe^-\bar{\nu}_e$
 - extraction of the g_{av}^Λ and g_w^Λ decay parameters
 - ♣ In progress
 - 4 If all steps work, consider more difficult scenario, mixed **MC** samples
 - 5 If previous step works, move to the real data
- Additional steps:
 - Sensitivity for g_i^Λ and A_i^Λ ($i = av, w$):
 - ♠ Preliminary result for $\Lambda \rightarrow pl\nu_l$ with $l = e, \mu$
 - $g_w^\Lambda = q^2 = 0$ - the same result for both decays
 - $g_w^\Lambda = 0$ and full q^2 range - similar result for both decays
 - $g_w^\Lambda \neq 0$ - better sensitivity for e -mode
 - Formalism with flip transition ($m_l^2/(2q^2) \neq 0$):
 - ♣ In progress

Backups




"I ALWAYS BACK UP EVERYTHING."


Random sample (e -mode, $N_{\text{sig}} = 10^5$) (step 1)




Random sample (e -mode, $N_{\text{phsp}} = 10^6$) (step 1)




Random sample (μ -mode, p , $N_{\text{sig}} = 10^5$) (step 1)




Random sample (μ -mode, p , $N_{\text{phsp}} = 10^6$) (step 1)



Random sample (μ -mode, $p \frac{(q^2 - m_\mu^2)^2}{q^2}$, $N_{\text{sig}} = 10^5$) (step 1)



Random sample (μ -mode, $p \frac{(q^2 - m_\mu^2)^2}{q^2}$, $N_{\text{phsp}} = 10^6$) (step 1)



Semileptonic Λ decay

- Momenta and masses: $\Lambda(p_1, M_1) \rightarrow p(p_2, M_2) + e^-(p_e, m_e) + \bar{\nu}_e(p_{\bar{\nu}_e}, 0)$
- Standard set of invariant FF for weak current-induced baryonic $\frac{1}{2}^+ \rightarrow \frac{1}{2}^+$ transitions:

$$M_\mu = M_\mu^V + M_\mu^A = \langle p(p_2) | J_\mu^{V+A} | \Lambda(p_1) \rangle =$$

$$= \bar{u}(p_2) \left[\gamma_\mu (F_1^V(q^2) + F_1^A(q^2) \gamma_5) + \frac{i\sigma_{\mu\nu} q^\nu}{M_1} (F_2^V(q^2) + F_2^A(q^2) \gamma_5) + \frac{q_\mu}{M_1} (F_3^V(q^2) + F_3^A(q^2) \gamma_5) \right] u(p_1)$$

where $q_\mu = (p_1 - p_2)_\mu$

- For $\Lambda \rightarrow p e^- \bar{\nu}_e$ at $\mathcal{O}(\frac{m_\pi^2}{2q^2}) \sim 4 \cdot 10^{-6} \Rightarrow F_3^{V,A} \rightarrow 0$
- $\Lambda \rightarrow p e^- \bar{\nu}_e \Rightarrow \Lambda \rightarrow p W^- (\rightarrow e^- \bar{\nu}_e)$
- Helicity amplitude is $H_{\lambda_2 \lambda_W} = (H_{\lambda_2 \lambda_W}^V + H_{\lambda_2 \lambda_W}^A)$ with $(\lambda_2 = \pm \frac{1}{2}; \lambda_W = 0, \pm 1)$:

$$\left. \begin{array}{l} \text{vector} \\ H_{\frac{1}{2}1}^V = \sqrt{2Q_-} \left(-F_1^V - \frac{M_1 + M_2}{M_1} F_2^V \right), \\ H_{\frac{1}{2}0}^V = \frac{\sqrt{Q_-}}{\sqrt{q^2}} \left((M_1 + M_2) F_1^V + \frac{q^2}{M_1} F_2^V \right), \end{array} \right\} \quad \left. \begin{array}{l} \text{axial-vector} \\ H_{\frac{1}{2}1}^A = \sqrt{2Q_+} \left(F_1^A - \frac{M_1 - M_2}{M_1} F_2^A \right), \\ H_{\frac{1}{2}0}^A = \frac{\sqrt{Q_+}}{\sqrt{q^2}} \left(-(M_1 - M_2) F_1^A + \frac{q^2}{M_1} F_2^A \right) \end{array} \right\}$$

$$\text{where } Q_\pm = (M_1 \pm M_2)^2 - q^2; \quad H_{-\lambda_2, -\lambda_W}^{V,A} = \pm H_{\lambda_2 \lambda_W}^{V,A}$$

Form factors

$$F_i^{V,A}(q^2) = F_i^{V,A}(0) \prod_{n=0}^{n_i} \frac{1}{1 - \frac{q^2}{m_{V,A}^2 + n\alpha'^{-1}}} \approx F_i^{V,A}(0) \left(1 + q^2 \sum_{n=0}^{n_i} \frac{1}{m_{V,A}^2 + n\alpha'^{-1}} \right)$$

	$F_i^{V,A}(0)(\Lambda \rightarrow p)$	$m_{V,A}$	$\alpha' [\text{GeV}^{-2}]$	n_i
$F_1^V(q^2)$	$-\sqrt{\frac{3}{2}}$ ¹	$m_{K^*(892)}^0$ ($J^P = 1^-$)	0.9	$n_1 = 1$
$F_2^V(q^2)$	$\frac{M_\Lambda \mu_p}{2M_p} F_1^V(0)$ ²			$n_2 = 2$
$F_3^V(q^2)$	0 ⁴			$n_3 = 2$
$F_1^A(q^2)$	$0.719 F_1^V(0)$ ³	$m_{K^*(1270)}^0$ ($J^P = 1^+$)		$n_1 = 1$
$F_2^A(q^2)$	0 ⁴	—		$n_2 = 2$
$F_3^A(q^2)$	$\frac{M_\Lambda(M_\Lambda + M_p)}{(m_K^-)^2} F_1^A(0)$ ⁴	m_K ($J^P = 0^-$)		$n_3 = 2$

● ¹ [PR135(1964)B1483], [PRL13(1964)264]


● ² $\mu_p = 1.793$ [Lect.NotesPhys.222(1985)1], [Ann.Rev.Nucl.Part.Sci.53(2003)39], [JHEP0807(2008)132]

● ³ [PRD41(1990)780]

● ⁴ Vanish in the $SU(3)$ symmetry limit; Goldberger-Treiman relation [PR110(1958)1178], [PR111(1958)354]

Decay chain: $\frac{1}{2} \rightarrow \frac{1}{2} + \{0, \pm 1\}$ (1)

- $e^+e^- \rightarrow (\Lambda \rightarrow pe^-\bar{\nu}_e)(\bar{\Lambda} \rightarrow \bar{p}\pi^+)$
- Formalism of $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ and $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ [PRD99(2019)056008]
- $\Lambda \rightarrow pe^-\bar{\nu}_e \Rightarrow \Lambda \rightarrow pW^- (\rightarrow e^-\bar{\nu}_e)$



- Decay matrix or transition matrix $b_{\mu\nu}$ for $\{\frac{1}{2} \rightarrow \frac{1}{2} + \{0, \pm 1\}\}$

$$b_{\mu\nu} = \frac{1}{4\pi} \sum_{\lambda_2=-1/2}^{1/2} \sum_{\lambda_W, \lambda'_W=-1}^1 H_{\lambda_2\lambda_W} H_{\lambda_2\lambda'_W}^* \sum_{\kappa, \kappa'=-1/2}^{1/2} (\sigma_\mu)^{\kappa, \kappa'} (\sigma_\nu)^{\lambda_2-\lambda'_W, \lambda_2-\lambda_W} \mathcal{D}_{\kappa, \lambda_2-\lambda_W}^{1/2*}(\Omega) \mathcal{D}_{\kappa', \lambda_2-\lambda'_W}^{1/2}(\Omega) \times$$

$$\sigma_\mu \rightarrow \sum_{\nu=0}^3 b_{\mu\nu} \sigma_\nu^{d-W} \Rightarrow \sum_{\lambda_l, \lambda_\nu=-1/2}^{1/2} |h'_{\lambda_l, \lambda_\nu=\pm 1/2}|^2 \mathcal{D}_{\lambda_W, \lambda_l-\lambda_\nu}^{1*}(\Omega') \mathcal{D}_{\lambda'_W, \lambda_l-\lambda_\nu}^1(\Omega'),$$

- Four helicity amplitudes: $H_{\frac{1}{2}0}, H_{-\frac{1}{2}0}, H_{\frac{1}{2}1}, H_{-\frac{1}{2}-1}$
- κ, κ' ; λ_2 - index of mother hyperon (Λ) and daughter baryon (p)
- λ_W, λ'_W ; λ_l, λ_ν - index of W^- -boson, lepton and neutrino
- **Kinematic variables:** $\Omega = \{\phi_p, \theta_p, 0\}$, $\Omega' = \{\chi, \theta_l, 0\}$ and $q^2 \in (m_e^2, (M_\Lambda - M_p)^2)$

Decay chain: $\frac{1}{2} \rightarrow \frac{1}{2} + \{0, \pm 1\}$ (2)

- Relation between helicity amplitudes and decay parameters

$$\begin{aligned}\sigma_D^{sl} &= \frac{1}{4}(1 \mp \cos \theta_l)^2 |H_{\frac{1}{2}1}|^2 + \frac{1}{4}(1 \pm \cos \theta_l)^2 |H_{-\frac{1}{2}-1}|^2 + \frac{1}{2} \sin^2 \theta_l (|H_{-\frac{1}{2}0}|^2 + |H_{\frac{1}{2}0}|^2), \\ \alpha_D^{sl} &= \frac{1}{4}(1 \mp \cos \theta_l)^2 |H_{\frac{1}{2}1}|^2 - \frac{1}{4}(1 \pm \cos \theta_l)^2 |H_{-\frac{1}{2}-1}|^2 + \frac{1}{2} \sin^2 \theta_l (|H_{-\frac{1}{2}0}|^2 - |H_{\frac{1}{2}0}|^2), \\ \beta_D^{sl} &= \frac{1}{\sqrt{2}} \sin \theta_l ((1 \pm \cos \theta_l) \Im(H_{-\frac{1}{2}0}^* H_{-\frac{1}{2}-1}) + (1 \mp \cos \theta_l) \Im(H_{\frac{1}{2}1}^* H_{\frac{1}{2}0})), \\ \gamma_D^{sl} &= \frac{1}{\sqrt{2}} \sin \theta_l ((1 \pm \cos \theta_l) \Re(H_{-\frac{1}{2}0}^* H_{-\frac{1}{2}-1}) + (1 \mp \cos \theta_l) \Re(H_{\frac{1}{2}1}^* H_{\frac{1}{2}0})).\end{aligned}$$

- Non-zero elements of the decay matrix $b_{\mu\nu}$:

$$\begin{aligned}b_{00} &= \sigma_D^{sl}, & b_{21} &= -(\gamma_D^{sl} \cos \chi + \beta_D^{sl} \sin \chi) \cos \theta_p \sin \phi_p \\ b_{03} &= \alpha_D^{sl}, & & -(\gamma_D^{sl} \sin \chi - \beta_D^{sl} \cos \chi) \cos \phi_p, \\ b_{10} &= \alpha_D^{sl} \cos \phi_p \sin \theta_p, & b_{22} &= (\gamma_D^{sl} \sin \chi - \beta_D^{sl} \cos \chi) \cos \theta_p \sin \phi_p \\ b_{11} &= -(\gamma_D^{sl} \cos \chi + \beta_D^{sl} \sin \chi) \cos \theta_p \cos \phi_p & & -(\gamma_D^{sl} \cos \chi + \beta_D^{sl} \sin \chi) \cos \phi_p, \\ &+ (\gamma_D^{sl} \sin \chi - \beta_D^{sl} \cos \chi) \sin \phi_p, & b_{23} &= \sigma_D^{sl} \sin \theta_p \sin \phi_p, \\ b_{12} &= (\gamma_D^{sl} \sin \chi - \beta_D^{sl} \cos \chi) \cos \theta_p \cos \phi_p & b_{30} &= \alpha_D^{sl} \cos \theta_p, \\ &+ (\gamma_D^{sl} \cos \chi + \beta_D^{sl} \sin \chi) \sin \phi_p, & b_{31} &= (\gamma_D^{sl} \cos \chi + \beta_D^{sl} \sin \chi) \sin \theta_p, \\ b_{13} &= \sigma_D^{sl} \sin \theta_p \cos \phi_p, & b_{32} &= -(\gamma_D^{sl} \sin \chi - \beta_D^{sl} \cos \chi) \sin \theta_p, \\ b_{20} &= \alpha_D^{sl} \sin \theta_p \sin \phi_p, & b_{33} &= \sigma_D^{sl} \cos \theta_p.\end{aligned}$$

- Main parameters:

$$\sigma_D^{sl} \equiv \sigma_D^{sl}(\theta_l, q^2), \quad \alpha_D^{sl} \equiv \alpha_D^{sl}(\theta_l, q^2), \quad \beta_D^{sl} \equiv \beta_D^{sl}(\theta_l, q^2), \quad \gamma_D^{sl} \equiv \gamma_D^{sl}(\theta_l, q^2)$$

- Each element of $b_{\mu\nu}$ is multiplied by $p = \sqrt{Q_+ Q_-} / (2M_1)$

Intermediate step (1)

- $\sigma_{\Lambda}^{sl}, \alpha_{\Lambda}^{sl}, \beta_{\Lambda}^{sl}$ and $\gamma_{\Lambda}^{sl} \Rightarrow \{n, \alpha, \alpha', \alpha'', \beta_{1,2}, \gamma_{1,2}\}(q^2): g_{av}^{\Lambda}(q^2), g_w^{\Lambda}(q^2)$
- Introduce the intermediate parameters:

$$\begin{aligned}
 \text{normalization} \quad n &= |H_{\frac{1}{2}1}|^2 + |H_{-\frac{1}{2}-1}|^2 + 2(|H_{-\frac{1}{2}0}|^2 + |H_{\frac{1}{2}0}|^2), \\
 \alpha &= |H_{\frac{1}{2}1}|^2 - |H_{-\frac{1}{2}-1}|^2 + 2(|H_{-\frac{1}{2}0}|^2 - |H_{\frac{1}{2}0}|^2), \\
 \alpha' &= |H_{\frac{1}{2}1}|^2 + |H_{-\frac{1}{2}-1}|^2 - 2(|H_{-\frac{1}{2}0}|^2 + |H_{\frac{1}{2}0}|^2), \\
 \alpha'' &= |H_{\frac{1}{2}1}|^2 - |H_{-\frac{1}{2}-1}|^2 - 2(|H_{-\frac{1}{2}0}|^2 - |H_{\frac{1}{2}0}|^2), \\
 \beta_{1,2} &= 2(\Im(H_{-\frac{1}{2}0}^* H_{-\frac{1}{2}-1}) \pm \Im(H_{\frac{1}{2}1}^* H_{\frac{1}{2}0})), \\
 \gamma_{1,2} &= 2(\Re(H_{-\frac{1}{2}0}^* H_{-\frac{1}{2}-1}) \pm \Re(H_{\frac{1}{2}1}^* H_{\frac{1}{2}0})),
 \end{aligned}$$

$$\text{where } \beta_{1,2} = \frac{1}{2}\sqrt{n^2 - \alpha^2 - (\alpha')^2 + (\alpha'')^2} \sin \phi_{1,2} \text{ and } \gamma_{1,2} = \frac{1}{2}\sqrt{n^2 - \alpha^2 - (\alpha')^2 + (\alpha'')^2} \cos \phi_{1,2}$$

- $\alpha^2 + (\alpha')^2 - (\alpha'')^2 + 2\sum_{i=1}^2(\gamma_i^2 + \beta_i^2) = n^2$
- Using the definition of helicity amplitudes the **main parameters** to describe semileptonic hyperon decays are:

$$\begin{aligned}
 &\bullet F_1^V(0), F_2^V(0), F_1^A(0) \\
 \text{or} \quad &\bullet g_{av}^D(0) = \frac{F_1^A(0)}{F_1^V(0)}, \quad g_w^D(0) = \frac{F_2^V(0)}{F_1^V(0)}
 \end{aligned}$$

Intermediate step (2)

- Relations between intermediate and decay parameters:

$$n = ((M_- M_+)^2 - q^4)(1 + (g_{av}^D(q^2))^2) + q^2 \left(4M_1 M_2 ((g_{av}^D(q^2))^2 - 1) + \frac{q^2}{M_1^2} Q_- ((g_w^D(q^2))^2 (M_+^2 + q^2) + 4g_w^D(q^2) M_+) \right)$$



$$\alpha = 2\sqrt{Q_- Q_+} \left[g_{av}^D(q^2)(q^2 - M_- M_+) + 2g_{av}^D(q^2)g_w^D(q^2)q^2 \frac{M_2}{M_1} \right]$$

$$\alpha' = Q_- Q_+ \left[-(1 + (g_{av}^D(q^2))^2) + (g_w^D(q^2))^2 \frac{q^2}{M_1^2} \right]$$

$$\alpha'' = 2\sqrt{Q_- Q_+} [g_{av}^D(q^2)(q^2 + M_- M_+) + 2g_{av}^D(q^2)g_w^D(q^2)q^2]$$

where $M_- = M_1 - M_2$ and $M_+ = M_1 + M_2$ and $Q_{\pm} = M_{\pm}^2 - q^2$

- $\{\alpha, \alpha', \alpha'', \gamma_{1,2}\}(q^2)/n(q^2) \in [-1, +1]$



Intermediate step (3)

- Non-zero elements of the decay matrix $b_{\mu\nu}$:

$$\begin{aligned}
 b_{00} &= 1, & b_{21} &= \mp A \cos \theta_p \sin \phi_p \mp B \cos \phi_p, \\
 b_{03} &= a_D^{sl}, & b_{22} &= \pm B \cos \theta_p \sin \phi_p \mp A \cos \phi_p, \\
 b_{10} &= a_D^{sl} \cos \phi_p \sin \theta_p, & b_{23} &= \sin \theta_p \sin \phi_p, \\
 b_{11} &= \mp A \cos \theta_p \cos \phi_p \pm B \sin \phi_p, & b_{30} &= a_D^{sl} \cos \theta_p, \\
 b_{12} &= \pm B \cos \theta_p \cos \phi_p \pm A \sin \phi_p, & b_{31} &= \pm A \sin \theta_p, \\
 b_{13} &= \sin \theta_p \cos \phi_p, & b_{32} &= \mp B \sin \theta_p, \\
 b_{20} &= a_D^{sl} \sin \theta_p \sin \phi_p, & b_{33} &= \cos \theta_p,
 \end{aligned}$$

$$\text{where } a_D^{sl} = \frac{\alpha_D^{sl}(\theta_l, q^2)}{\sigma_D^{sl}(\theta_l, q^2)} = \frac{\alpha + \alpha'' \cos^2 \theta_l \mp (n + \alpha') \cos \theta_l}{n + \alpha' \cos^2 \theta_l \mp (\alpha + \alpha'') \cos \theta_l},$$

$$A = \frac{1}{2\sqrt{2}} \frac{\sin \theta_l}{\sigma_D^{sl}(\theta_l, q^2)} [\cos \chi(\gamma_1 \pm \cos \theta_l \gamma_2) + \sin \chi(\beta_1 \pm \cos \theta_l \beta_2)],$$

$$B = \frac{1}{2\sqrt{2}} \frac{\sin \theta_l}{\sigma_D^{sl}(\theta_l, q^2)} [\sin \chi(\gamma_1 \pm \cos \theta_l \gamma_2) - \cos \chi(\beta_1 \pm \cos \theta_l \beta_2)].$$

Helicity amplitudes of the lepton pair $h_{\lambda_l \lambda_\nu}^l$

- Lepton and anti-neutrino spinors

$$\bar{u}_l(\mp \frac{1}{2}, p_{l-}) = \sqrt{E_l + m_l} \left(\chi_{\mp}^{\dagger}, \frac{\pm |\vec{p}_{l-}|}{E_l + m_l} \chi_{\mp}^{\dagger} \right),$$

$$v_{\bar{\nu}}(\frac{1}{2}) = \sqrt{E_{\nu}} \begin{pmatrix} \chi_+ \\ -\chi_+ \end{pmatrix},$$

where $\chi_+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\chi_- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are Pauli two-spinors

- SM form of the lepton current ($\lambda_W = \lambda_{l-} - \lambda_{\bar{\nu}}$)

$$h_{\lambda_{l-}=\mp 1/2, \lambda_{\bar{\nu}}=1/2}^l = \bar{u}_l(\mp \frac{1}{2}) \gamma^{\mu} (1 + \gamma_5) v_{\bar{\nu}}(\frac{1}{2}) \left\{ \begin{matrix} \epsilon_{\mu}(-1) \\ \epsilon_{\mu}(t), \epsilon_{\mu}(0) \end{matrix} \right\}$$

where $\epsilon^{\mu}(t) = (1; 0, 0, 0)$, $\epsilon^{\mu}(0) = (0; 0, 0, 1)$ and $\epsilon^{\mu}(\mp 1) = (0; \mp 1, -i, 0)/\sqrt{2}$

- Moduli squared of the helicity amplitudes [\[EPJ C59 \(2009\) 27\]](#)

$$\text{nonflip}(\lambda_W = \mp 1) : |h_{\lambda_l=\mp \frac{1}{2}, \lambda_{\nu}=\pm \frac{1}{2}}^l|^2 = 8(q^2 - m_l^2),$$

$$\text{flip}(\lambda_W = 0, t) : |h_{\lambda_l=\pm \frac{1}{2}, \lambda_{\nu}=\pm \frac{1}{2}}^l|^2 = 8 \frac{m_l^2}{2q^2} (q^2 - m_l^2)$$

- Upper and lower signs refer to the configurations $(l^-, \bar{\nu}_l)$ ($\lambda_{\nu} = 1/2$) and (l^+, ν_l) ($\lambda_{\nu} = -1/2$), respectively
- In case of [the e-mode](#) only [nonflip transition](#) remains under assumption $\frac{m_e^2}{2q^2} \rightarrow 0$

Semileptonic Λ decay^[flip]

- For $\Lambda \rightarrow pe^- \bar{\nu}_e$ at $\mathcal{O}(\frac{m_e^2}{2q^2}) \sim 4 \cdot 10^{-6} \Rightarrow F_3^{V,A} \rightarrow 0$
- For $\Lambda \rightarrow p\mu^- \bar{\nu}_\mu$ at $\mathcal{O}(\frac{m_\mu^2}{2q^2}) \sim 0.18 \Rightarrow F_3^{V,A} \neq 0$
- Helicity amplitude is $H_{\lambda_2 \lambda_W} = (H_{\lambda_2 \lambda_W}^V + H_{\lambda_2 \lambda_W}^A)$ with $(\lambda_2 = \pm \frac{1}{2}; \lambda_W = t, 0, \pm 1)$:

$$\begin{array}{c} \text{vector} \\ \left[\begin{array}{l} H_{\frac{1}{2}1}^V = \sqrt{2Q_-} \left(-F_1^V - \frac{M_1 + M_2}{M_1} F_2^V \right), \\ H_{\frac{1}{2}0}^V = \frac{\sqrt{Q_-}}{\sqrt{q^2}} \left((M_1 + M_2) F_1^V + \frac{q^2}{M_1} F_2^V \right), \\ H_{\frac{1}{2}t}^V = \frac{\sqrt{Q_+}}{\sqrt{q^2}} \left((M_1 - M_2) F_1^V + \frac{q^2}{M_1} F_3^V \right), \end{array} \right. \end{array} \quad \begin{array}{c} \text{axial-vector} \\ \left[\begin{array}{l} H_{\frac{1}{2}1}^A = \sqrt{2Q_+} \left(F_1^A - \frac{M_1 - M_2}{M_1} F_2^A \right), \\ H_{\frac{1}{2}0}^A = \frac{\sqrt{Q_+}}{\sqrt{q^2}} \left(-(M_1 - M_2) F_1^A + \frac{q^2}{M_1} F_2^A \right), \\ H_{\frac{1}{2}t}^A = \frac{\sqrt{Q_-}}{\sqrt{q^2}} \left(-(M_1 + M_2) F_1^A + \frac{q^2}{M_1} F_3^A \right), \end{array} \right. \end{array}$$

$$\text{where } Q_{\pm} = (M_1 \pm M_2)^2 - q^2; \quad H_{-\lambda_2, -\lambda_W}^{V,A} = \pm H_{\lambda_2 \lambda_W}^{V,A}$$

- SU(3) symmetry limit $\Rightarrow F_3^V = 0$

Decay chain: $\frac{1}{2} \rightarrow \frac{1}{2} + \{t, 0, \pm 1\}$ [flip]

- Six helicity amplitudes: $H_{\frac{1}{2}0}, H_{-\frac{1}{2}0}, H_{\frac{1}{2}1}, H_{-\frac{1}{2}-1}, H_{\frac{1}{2}t}, H_{-\frac{1}{2}t}$
- Relation between helicity amplitudes and decay parameters

$$\begin{aligned}\sigma_D'^{sl} &= \frac{1}{2} \sin^2 \theta_l (|H_{\frac{1}{2}1}|^2 + |H_{-\frac{1}{2}-1}|^2) + \cos^2 \theta_l (|H_{-\frac{1}{2}0}|^2 + |H_{\frac{1}{2}0}|^2) + |H_{-\frac{1}{2}t}|^2 + |H_{\frac{1}{2}t}|^2 - 2 \cos \theta_l \Re(H_{-\frac{1}{2}t} H_{-\frac{1}{2}0}^* + H_{\frac{1}{2}t} H_{\frac{1}{2}0}^*), \\ \alpha_D'^{sl} &= \frac{1}{2} \sin^2 \theta_l (|H_{\frac{1}{2}1}|^2 - |H_{-\frac{1}{2}-1}|^2) + \cos^2 \theta_l (|H_{-\frac{1}{2}0}|^2 - |H_{\frac{1}{2}0}|^2) + |H_{-\frac{1}{2}t}|^2 - |H_{\frac{1}{2}t}|^2 - 2 \cos \theta_l \Re(H_{-\frac{1}{2}t} H_{-\frac{1}{2}0}^* - H_{\frac{1}{2}t} H_{\frac{1}{2}0}^*), \\ \beta_{3,4} &= 2\Im(H_{-\frac{1}{2}t}^* H_{-\frac{1}{2}-1} \pm H_{\frac{1}{2}1}^* H_{\frac{1}{2}t}), \\ \gamma_{3,4} &= 2\Re(H_{-\frac{1}{2}t}^* H_{-\frac{1}{2}-1} \pm H_{\frac{1}{2}1}^* H_{\frac{1}{2}t}).\end{aligned}$$

- Non-zero elements of the decay matrix $b'_{\mu\nu}$:

$$\begin{aligned}b'_{00} &= 1, & b'_{21} &= B' \cos \phi_p + A' \sin \phi_p \cos \theta_p, \\ b'_{03} &= a_D'^{sl}, & b'_{22} &= A' \cos \phi_p - B' \sin \phi_p \cos \theta_p, \\ b'_{10} &= a_D'^{sl} \cos \phi_p \sin \theta_p, & b'_{23} &= \sin \theta_p \sin \phi_p, \\ b'_{11} &= A' \cos \theta_p \cos \phi_p - B' \sin \phi_p, & b'_{30} &= a_D'^{sl} \cos \theta_p, \\ b'_{12} &= -B' \cos \theta_p \cos \phi_p - A' \sin \phi_p, & b'_{31} &= -A' \sin \theta_p, \\ b'_{13} &= \sin \theta_p \cos \phi_p, & b'_{32} &= B' \sin \theta_p, \\ b'_{20} &= a_D'^{sl} \sin \theta_p \sin \phi_p, & b'_{33} &= \cos \theta_p, \\ \text{where } a_D'^{sl} &= \frac{\alpha_D'^{sl}(\theta_l, q^2)}{\sigma_D'^{sl}(\theta_l, q^2)}, & A' &= \frac{1}{\sqrt{2} \sigma_D'^{sl}(\theta_l, q^2)} [\cos \chi (\cos \theta_l \gamma_2 - \gamma_4) + \sin \chi (\cos \theta_l \beta_2 - \beta_4)], \\ & & B' &= \frac{1}{\sqrt{2} \sigma_D'^{sl}(\theta_l, q^2)} [\sin \chi (\cos \theta_l \gamma_2 - \gamma_4) - \cos \chi (\cos \theta_l \beta_2 - \beta_4)].\end{aligned}$$

- Each element of $b'_{\mu\nu}$ is multiplied by $p = \sqrt{Q_+ Q_-} / (2M_1)$ and $\frac{m_l^2}{2q^2}$

Intermediate step^[flip] (1)

- $\sigma_{\Lambda}^{'sl}$ and $\alpha_{\Lambda}^{'sl} \Rightarrow \{n, \alpha, \alpha', \alpha'', \beta_{1,2}, \gamma_{1,2}\}(q^2) + \{\epsilon, \epsilon', \beta_{3,4}, \gamma_{3,4}\}(q^2): g_{av}^{\Lambda}(q^2), g_{av3}^{\Lambda}(q^2)$
- Additional intermediate parameters:

$$\epsilon = 4(|H_{-\frac{1}{2}t}|^2 + |H_{\frac{1}{2}t}|^2),$$

$$\epsilon' = 4(|H_{-\frac{1}{2}t}|^2 - |H_{\frac{1}{2}t}|^2).$$

- $(n + \alpha')(n - \alpha' + \epsilon) - (\alpha + \alpha'')(\alpha - \alpha'' + \epsilon') - 2 \sum_{i=1}^4 (\gamma_i^2 + \beta_i^2) = 0$
If no flip transition $\Rightarrow$ relation on slide 22
- Introduce additional main parameter of semileptonic hyperon decay:

$$\bullet F_3^A(0)$$

or

$$\bullet g_{av3}^D(0) = \frac{F_3^A(0)}{F_1^V(0)}$$

Intermediate step^[nonflip+flip] (2)

- Relations between intermediate and decay parameters:

$$n = ((M_- M_+)^2 - q^4)(1 + (g_{av}^D(q^2))^2) + q^2 \left(4M_1 M_2 ((g_{av}^D(q^2))^2 - 1) + \frac{q^2}{M_1^2} Q_- ((g_w^D(q^2))^2 (M_+^2 + q^2) + 4g_w^D(q^2) M_+) \right)$$

$$\alpha = 2\sqrt{Q_- Q_+} \left[g_{av}^D(q^2)(q^2 - M_- M_+) + 2g_{av}^D(q^2)g_w^D(q^2)q^2 \frac{M_2}{M_1} \right]$$

$$\alpha' = Q_- Q_+ \left[-(1 + (g_{av}^D(q^2))^2) + (g_w^D(q^2))^2 \frac{q^2}{M_1^2} \right]$$



$$\alpha'' = 2\sqrt{Q_- Q_+} [g_{av}^D(q^2)(q^2 + M_- M_+) + 2g_{av}^D(q^2)g_w^D(q^2)q^2]$$

$$\epsilon = (M_- M_+)^2 (1 + (g_{av}^D(q^2))^2) - q^2 \left(M_-^2 + M_+^2 (g_{av}^D(q^2))^2 + \frac{Q_-}{M_1} (2g_{av}^D(q^2)g_{av3}^D(q^2)M_+ - (g_{av3}^D(q^2))^2 q^2) \right)$$

$$\epsilon' = 2M_- \sqrt{Q_- Q_+} \left(g_{av}^D(q^2)M_+ - \frac{q^2}{M_1} g_{av3}^D(q^2) \right)$$

where $M_- = M_1 - M_2$ and $M_+ = M_1 + M_2$ and $Q_{\pm} = M_{\pm}^2 - q^2$

- $\{\alpha, \alpha', \alpha'', \gamma_{1,2}\}(q^2) + \{\epsilon, \epsilon', \gamma_{3,4}\}(q^2)/n(q^2) \in [-1, +1]$



Joint angular distribution

- Full decay matrix of semileptonic hyperon decay:

$$b_{\mu\nu}^f = \sum_{\mu,\nu=0}^3 \left(b_{\mu\nu} + \frac{m_l^2}{2q^2} b'_{\mu\nu} \right)$$

where $b_{\mu\nu}$ and $b'_{\mu\nu}$ are nonflip (slide 24) and flip (slide 27) transitions

- Process $e^+e^- \rightarrow (\Lambda \rightarrow p e^- \bar{\nu}_e)(\bar{\Lambda} \rightarrow \bar{p} \pi^+)$

$$\text{Tr} \rho_{pW\bar{p}} \propto \mathcal{W}(\xi; \omega) = \sum_{\mu, \bar{\nu}=0}^3 C_{\mu\bar{\nu}} b_{\mu 0}^{\Lambda} a_{\bar{\nu} 0}^{\bar{\Lambda}}$$


- $C_{\mu\bar{\nu}}(\theta_{\Lambda}; \alpha_{\psi}, \Delta\Phi)$
- $b_{\mu 0}^f$ matrices for $1/2 \rightarrow 1/2 + \{t, 0, \pm 1\}$ decays $\Leftrightarrow b_{\mu 0}^{\Lambda} \equiv b_{\mu 0}^f(\theta_p, \varphi_p, \theta_e, \chi, q^2; g_{av}^{\Lambda}, g_w^{\Lambda}, g_{av3}^{\Lambda})$
- $a_{\bar{\nu} 0}$ matrices for $1/2 \rightarrow 1/2 + 0$ decays $\Leftrightarrow a_{\bar{\nu} 0}^{\bar{\Lambda}} \equiv a_{\bar{\nu} 0}(\theta_{\bar{p}}, \varphi_{\bar{p}}, \alpha_{\bar{\Lambda}})$
 - $\xi \equiv (\theta_{\Lambda}, \theta_p, \varphi_p, \theta_e, \chi, q^2, \theta_{\bar{p}}, \varphi_{\bar{p}})$
 - $\omega \equiv (\alpha_{\psi}, \Delta\Phi, g_{av}^{\Lambda}, g_w^{\Lambda}, g_{av3}^{\Lambda}, \alpha_{\bar{\Lambda}})$
- If flip transition is taking into account, $g_{av3}^D \neq 0$
- Range of $q^2 \in (m_l^2, (M_1 - M_2)^2)$ is specific for each decay

Size estimations of helicity amplitudes

$$T(V,A) := \sqrt{q^2} H_{\frac{1}{2}1}^{V,A}$$

$$L(V,A) := \sqrt{q^2} H_{\frac{1}{2}0}^{V,A}$$

$$S(V,A) := \sqrt{q^2} H_{\frac{1}{2}t}^{V,A}$$



Dashed lines:
 $F_2^V(q^2)$ and $F_3^A(q^2)$
 are switched off

- If $q^2 = m_e^2 \Rightarrow H_{\frac{1}{2}0}^V \sim H_{\frac{1}{2}t}^V$ and $H_{\frac{1}{2}0}^A \sim H_{\frac{1}{2}t}^A$ are dominated
- If $q^2 = (M_\Lambda - M_p)^2 \Rightarrow H_{\frac{1}{2}t}^V$ and $H_{\frac{1}{2}1}^A = -\sqrt{2}H_{\frac{1}{2}0}^A$ are dominated
- Using data of the E-555 experiment (Fermilab) [\[PRD41 \(1990\) 780\]](#)

$$\Rightarrow \frac{F_1^A(0)}{F_1^V(0)} (\Lambda \rightarrow pe^- \bar{\nu}_e) = 0.731 \pm 0.016 \text{ and } \frac{F_2^V(0)}{F_1^V(0)} (\Lambda \rightarrow pe^- \bar{\nu}_e) = 0.15 \pm 0.30$$

$$\Rightarrow \frac{F_1^A(0)}{F_1^V(0)} (\Lambda \rightarrow pe^- \bar{\nu}_e) = 0.719 \pm 0.016 \text{ with constraint } \frac{F_2^V(0)}{F_1^V(0)} (\Lambda \rightarrow pe^- \bar{\nu}_e) \rightarrow 0.97 \text{ (CVC)}$$

$$A_{\Lambda} = \frac{\alpha_{\Lambda} + \alpha_{\bar{\Lambda}}}{\alpha_{\Lambda} - \alpha_{\bar{\Lambda}}}$$

- BESIII result: $A_{\Lambda} = -0.006 \pm 0.012 \pm 0.007$ [NaturePhys.15(2019)631]
 - CKM: $-3 \cdot 10^{-5} \leq A_{\Lambda} \leq 4 \cdot 10^{-5}$ [PRD67(2003)056001]
- Extensions of SM: $A_{\Lambda \rightarrow p\pi^-} \sim (0.05 - 1.2) \cdot 10^{-4}$ [Chin.Phys.C42(2018)013101]

Experiment	N _{evt}	$\sigma(A_{\Lambda})$	
BESIII (2018)	$4.2 \cdot 10^5$	$1.2 \cdot 10^{-2}$	$N_{J/\psi} = 1.31 \cdot 10^9$
Some estimations			
BESIII	$3 \cdot 10^6$	$5 \cdot 10^{-3}$	$N_{J/\psi} = 10^{10}$ $\mathcal{L} = 0.47 \cdot 10^{33}/\text{cm}^2/\text{s}$, $\Delta E = 0.9 \text{ MeV}$
SuperTauCharm	$6 \cdot 10^8$	$3 \cdot 10^{-4}$	$N_{J/\psi} = 2 \cdot 10^{12}$ $\mathcal{L} = 10^{35}/\text{cm}^2/\text{s}$, $\Delta E = 0.9 \text{ MeV}$
SuperTauCharm + reduced ΔE	$3 \cdot 10^9$	$1.4 \cdot 10^{-4}$	$N_{J/\psi} = 10^{13}$ $\mathcal{L} = 10^{35}/\text{cm}^2/\text{s}$, $\Delta E < 0.9 \text{ MeV}$ (?)

CP symmetry test A_{Λ}^{av}

$$A_{\Lambda}^{av} = \frac{g_{av}^{\Lambda}(0) + g_{av}^{\bar{\Lambda}}(0)}{g_{av}^{\Lambda}(0) - g_{av}^{\bar{\Lambda}}(0)}$$

- $g_{av}(0) = \frac{F_1^A(0)}{F_1^V(0)}$
 - Determination of $F_1^V(0)$ value by CVC hypothesis: $F_1^V(0) < 0$
 - No determination of $F_1^A(0)$ value by any general theoretical arguments
- Some assumptions:
 - $F_1^A(0) \leq 0$ then $g_{av}(0) \geq 0$
 - ? Naïve assumption $g_{av}^{\Lambda} = -g_{av}^{\bar{\Lambda}}$
 $\Rightarrow$ Need to be care with sign of g_{av}

Experiment	N _{evt}	Result	Reference
E555 (Fermilab)	37286	$g_{av}(0) = 0.719 \pm 0.016 \pm 0.012$ constrain on $g_w(0) = 0.97$	[PRD41(1990)780]
SPS (CERN)	7111	$g_{av}(0) = 0.70 \pm 0.03$ measured $F_2^V(0) = 1.32 \pm 0.81$	[ZPC21(1983)1]
AGS (BNL)	10 ⁴	$ g_{av}(0) = 0.734 \pm 0.031$ used $g_w(0) = 0.97$	[PLB98(1981)123]

Boundary case: $q_{\min}^2$

- $q_{\min}^2 = m_e^2 \longrightarrow 0$:

$$b_{00} = (1 + (g_{av}^D(0))^2) \sin^2 \theta_l,$$

$$b_{03} = -2g_{av}^D(0) \sin^2 \theta_l,$$

$$b_{10} = b_{03} \sin \theta_p \cos \phi_p,$$

$$b_{13} = b_{00} \sin \theta_p \cos \phi_p,$$

$$b_{20} = b_{03} \sin \theta_p \sin \phi_p,$$

$$b_{23} = b_{00} \sin \theta_p \sin \phi_p,$$

$$b_{30} = b_{03} \cos \theta_p,$$

$$b_{33} = b_{00} \cos \theta_p$$